herpy

LEGWANY

Hernandez's Helmeted Basilisk

Corytophanes hernandesii

Corytophanes hernandesii is a slender, arboreal lizard from the forests of southern Mexico, Belize, Guatemala and Honduras with an angular, bony helmet on the head. It sits motionless for hours waiting for large insects, changes colour from brown to yellow, orange and almost black, and is very shy. It needs a tall, humid enclosure, UVB, large live feeder insects and peace.

Rodzina: Corytophanidae Dawna nazwa: *Chamaeleopsis hernandesii; Corytophanes chamaeleopsis; Corytophanes hernandezii; Corytophanes mexicanus*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**25-35 cm** długość całkowita z ogonem
Długość życia	**5-8 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny**
Środowisko	**Suchy las i las deszczowy**
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**60 × 45 × 90 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**26-30 °C**
Miejsce wygrzewania	**32-35 °C**
Temperatura w nocy	**21-24 °C**
Wilgotność	**60-90%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 1
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub jeden samiec z kilkoma samicami**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**3-11 jaj**
Inkubacja	**90 dni** przy 26 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	A single adult needs a tall enclosure of at least 60 × 45 × 90 cm (length × width × height). Larger is better, especially wider.
TEMPERATURA	The air should be 26-30 °C by day, and the warmest spot, a thick branch under a lamp, should have a surface temperature of 32-35 °C, measured with an infrared thermometer.
WILGOTNOŚĆ I WODA	Humidity should be 60-90 %, lowest in the middle of the day and highest after misting and at night.
OŚWIETLENIE I UVB	*Corytophanes hernandesii* is diurnal and needs UVB all day, even though it is a shade species. The UVB lamp should give Ferguson zone 1, i.e. a low UV index on the highest branches and almost none lower in the enclosure.
ŻYWIENIE	The species is insectivorous and takes large, slow prey: cockroaches (such as discoids and dubia), silkworms, hornworms, grasshoppers, superworms and earthworms.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is shy and peaceful but sensitive to disturbance, and it is better suited to observing than to handling. It often sits completely still for hours and moves slowly.
NATURALNY ZASIĘG	Meksyk, Belize, Gwatemala i Honduras.