herpy

LEGWANY

Desert horned lizard

Phrynosoma platyrhinos

Phrynosoma platyrhinos is a broad, flattened horned lizard from the deserts of the western USA and north-western Mexico. It reaches 7-10 cm from snout to vent, is diurnal and burrows into loose sand. It feeds mostly on ants, and it is precisely this diet that makes it one of the most difficult lizards to keep in an enclosure. There is little knowledge of how it should be cared for.

Rodzina: Phrynosomatidae Dawna nazwa: *Anota calidiarum; Doliosaurus platyrhinos*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**7-10 cm** długość od pyska do kloaki
Długość życia	**7-9 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny i kopiący nory**
Środowisko	**Pustynia i półpustynia**
Trudność	**Ekspert**

HODOWLA

Min. terrarium	**100 × 50 × 40 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**27-30 °C**
Miejsce wygrzewania	**38-42 °C**
Temperatura w nocy	**18-22 °C**
Wilgotność	**20-40%**
Woda	**Bez miski z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady**
Utrzymanie	**Pojedynczo**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**2-16 jaj**
Zniesienia w roku	**1-2**
Inkubacja	**40-61 dni** przy 27-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	There is little knowledge about the care of *Phrynosoma platyrhinos*. Care is therefore based on the habitat: open, hot sandy flats, loose substrate, strong sun and a diet of ants.
TEMPERATURA	The species is a sun lover and needs a basking spot where the surface reaches 38-42 °C, measured with an infrared thermometer.
WILGOTNOŚĆ I WODA	Humidity should be low, 20-40%, and the enclosure should dry out completely during the day. The substrate must not be wet. The species hardly ever drinks from a dish, but licks droplets from stones and plants.
OŚWIETLENIE I UVB	The species is diurnal and spends the day in open sunlight, so it needs strong light and UVB all year. Use a UVB tube (Ferguson zone 3, T5 HO 10-14%) or a combined UVB metal halide lamp over the basking spot.
ŻYWIENIE	Food is the decisive limitation. In the wild, ants make up roughly half to four fifths of the diet, and the species eats large quantities. The rest is beetles, crickets, grasshoppers and flies.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Phrynosoma platyrhinos* is calm and lives alone. It does not make much fuss and is easy to overlook, because it blends into the ground. It moves in short bursts and stops again.
NATURALNY ZASIĘG	Stany Zjednoczone i Meksyk.