herpy

SALAMANDRY I TRASZKI

Marbled Salamander

Ambystoma opacum

Ambystoma opacum is a small, stocky salamander of 9-11 cm from eastern North America, black with crossbands of white or silvery grey. The species has an unusual breeding biology: mating and egg-laying take place on land in autumn, and the female stays with the eggs until rain fills the hollow and the larvae hatch. The adults live hidden and do best kept cool.

Rodzina: Ambystomatidae Dawna nazwa: *Salamandra armigera; Salamandra fasciata* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**9-11 cm** długość całkowita z ogonem
Długość życia	**4-10 lat**
Aktywność	**W nocy**
Tryb życia	**Naziemny i kopiący nory**
Środowisko	**Lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Początkujący**

HODOWLA

Min. terrarium	**40 × 25 × 25 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**18-24 °C**
Temperatura w nocy	**16-20 °C**
Wilgotność	**80-95%**
Woda	**Miska z wodą** głębokość 1-2 cm
UVB	**Niewymagane**
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**Na ziemi**
Dojrzałość płciowa	**12-24 miesięcy**
Liczba jaj w zniesieniu	**50-200**
Zniesienia w roku	**1**
Okres larwalny	**60-270 dni** od wyklucia do metamorfozy
Pokarm larw	**Inne larwy i małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Pilnuje jaj**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	There is little knowledge about the husbandry of *Ambystoma opacum*, and the advice on substrate and water is partly borrowed from the closely related *Ambystoma maculatum* and *Ambystoma tigrinum*, which are kept far more often.
TEMPERATURA	By day the temperature should be 18-24 °C, and at night it can be a couple of degrees cooler, 16-20 °C. The species tolerates cold better than heat, and above 30 °C for long periods it can die.
WILGOTNOŚĆ I ZRASZANIE	The substrate must be damp, but never wet or muddy, because the animal absorbs water through its skin. A damp substrate corresponds to 80-95% humidity. Mist the substrate with chlorine-free water when the surface looks dry.
WODA I JAKOŚĆ WODY	Adults are poor swimmers, and the species drowns if it cannot get out of the water. Deep dishes and water sections must therefore be avoided.
OŚWIETLENIE I UVB	The species is nocturnal and lives hidden, so it does not need special lighting and does not need UVB either. UV light can damage the eyes and skin of amphibians, and larvae must not be exposed to it.
ŻYWIENIE	Adults eat crawling and flying insects, worms and small snails: crickets, earthworms and compost worms, and occasionally waxworms. The feeder animals must be smaller than the animal's head.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	Adults lie hidden most of the time and come out at night to hunt. The species does not bite humans. Outside the breeding season the animals live alone, and males in particular bite each other on the snout if they meet.
NATURALNY ZASIĘG	Stany Zjednoczone.