herpy

SALAMANDRY I TRASZKI

Anderson's Crocodile Newt

Echinotriton andersoni

Echinotriton andersoni is a compact, dark brown to black salamander, 14-21 cm long, from the central Ryukyu Islands of Japan. It lives mostly on land in damp forest, has knob-like warts along its sides and can push the tips of its ribs out through the skin as a defence. The species is rare and little known in captivity.

Rodzina: Salamandridae Dawna nazwa: *Pleurodeles andersoni; Tylototriton andersoni* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**14-21 cm** długość całkowita z ogonem
Długość życia	**10-20 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Naziemny**
Środowisko	**Suchy las i mokradła**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**60 × 40 × 40 cm** D × S × W dla 3 dorosłych osobników
Temperatura w dzień	**18-25 °C**
Temperatura w nocy	**15-20 °C**
Wilgotność	**60-80%**
Woda	**Część wodna** głębokość 10-20 cm
UVB	**Niewymagane**
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**Na ziemi**
Liczba jaj w zniesieniu	**3-46**
Czas do wyklucia	**20-27 dni**
Okres larwalny	**70-100 dni** od wyklucia do metamorfozy
Pokarm larw	**Inne larwy i małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik III**
Załącznik unijny	**Załącznik C**
IUCN	**Narażony (VU)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Echinotriton andersoni*.
TEMPERATURA	The temperature is 18-25 °C by day and may fall to 15-20 °C at night. It should not stay above 25 °C for long. Heat lamps are unnecessary; keep the terrarium cool in summer and out of direct sun. Winter rest is not documented.
WILGOTNOŚĆ I ZRASZANIE	Humidity should be 60-80%. The leaf litter and soil are kept damp but not wet, and part of the substrate should be drier so the animals can choose. Provide a little air exchange so the air does not stand still.
WODA I JAKOŚĆ WODY	The water area is a bowl-shaped depression, about 30 cm in diameter and 10-20 cm deep in the middle, with an even slope all the way round, so the animals are never at risk of drowning.
OŚWIETLENIE I UVB	The species lives hidden and is most active at dusk and at night. UVB is not necessary. A light period of about 10 hours is suitable, and the light must also be sufficient to keep the plants alive. Avoid direct sun.
ŻYWIENIE	The animals take earthworms, crickets, snails, woodlice and small beetles. Earthworms and crickets are suitable as the staple, and the feeder insects must be small enough for the animal to swallow whole.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The animal lives hidden and is most visible at dusk and at night. Groups need plenty of hiding places. The animal is not suited to handling.
NATURALNY ZASIĘG	Japonia.