herpy

SALAMANDRY I TRASZKI

Chinhai Spiny Newt

Echinotriton chinhaiensis

Echinotriton chinhaiensis is a critically endangered salamander found only in a small area of Beilun District, east of Ningbo in Zhejiang, China. The eggs are laid on land beside small, still ponds, and the larvae live in the water for 58-88 days. The species reaches sexual maturity only after about ten years and can live to at least 20 years. Almost nothing is known about the care of adults.

Rodzina: Salamandridae Dawna nazwa: *Pleurodeles chinhaiensis; Tylototriton chinhaiensis* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**14-21 cm** długość całkowita z ogonem
Długość życia	**20-25 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Naziemny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**60 × 40 × 40 cm** D × S × W dla 3 dorosłych osobników
Temperatura w dzień	**18-25 °C**
Temperatura w nocy	**15-20 °C**
Wilgotność	**60-80%**
Woda	**Część wodna** głębokość 10-20 cm
UVB	**Niewymagane**
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**Na ziemi**
Liczba jaj w zniesieniu	**3-46**
Czas do wyklucia	**29-31 dni** przy 20 °C
Okres larwalny	**58-88 dni** od wyklucia do metamorfozy
Pokarm larw	**Inne larwy i małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Krytycznie zagrożony (CR)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Echinotriton chinhaiensis*. Most studies concern eggs and larvae.
TEMPERATURA	The temperature for adults is borrowed from *Echinotriton andersoni*: 18-25 °C by day and 15-20 °C at night. The requirements of adults have not been studied in *Echinotriton chinhaiensis*.
WILGOTNOŚĆ I ZRASZANIE	Humidity is borrowed from *Echinotriton andersoni*: 60-80 %. Keep the leaf litter and soil damp but not wet, and part of the substrate should be drier, so the animals can choose. Eggs on land must be kept moist at all times.
WODA I JAKOŚĆ WODY	The water area is a bowl-shaped depression, about 30 cm across and 10-20 cm deep in the middle, with an even slope all the way round, so the animals are not at risk of drowning.
OŚWIETLENIE I UVB	UVB is not necessary. The animals are assumed to be most active at dusk and at night; this is borrowed from *Echinotriton andersoni*. A light period of about 10 hours suits, and the light must also be able to keep the plants alive.
ŻYWIENIE	The diet of adults is not documented. In *Echinotriton andersoni* it is earthworms, crickets, snails, woodlice and small beetles, and the food must be small enough to be swallowed whole.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The behaviour of adults is not documented. The larvae are cautious: they avoid visible predators and reduce their activity when they sense predators by smell or sight. The animals are not suited to handling.
NATURALNY ZASIĘG	Chiny.