herpy

SALAMANDRY I TRASZKI

Bosca's newt

Lissotriton boscai

Lissotriton boscai is a small, mostly aquatic newt of 7-9 cm from Portugal and western Spain. It is brownish to greenish with dark spots and an orange belly, and in the breeding season the male develops a coloured protuberance on the tip of the tail instead of a crest. The species is cool-loving and adaptable and thrives in a cool aquaterrarium.

Rodzina: Salamandridae Dawna nazwa: *Triturus boscai* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**7-9 cm** długość całkowita z ogonem
Długość życia	**4-12 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Półwodny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**60 × 30 × 30 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**12-20 °C**
Temperatura w nocy	**8-15 °C**
Wilgotność	**80-100%**
Woda	**Akwaterrarium** głębokość 10-20 cm
UVB	**Niewymagane**
Zimowanie	**6-10 °C** 6-8 tygodni
Pokarm	**Owady, małe bezkręgowce (skoczogonki, muszki owocowe) i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**W wodzie**
Dojrzałość płciowa	**24-36 miesięcy**
Liczba jaj w zniesieniu	**150-440**
Czas do wyklucia	**8-28 dni**
Okres larwalny	**45-105 dni** od wyklucia do metamorfozy
Pokarm larw	**Małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	*Lissotriton boscai* is mostly aquatic, and an aquaterrarium suits it best. An enclosure of 60 × 30 × 30 cm holds up to 4 animals, and a larger one gives more peace.
TEMPERATURA	*Lissotriton boscai* is cool-loving. The daytime temperature is 12-20 °C and the night-time temperature 8-15 °C. Temperatures above 22 °C for long periods should be avoided.
WILGOTNOŚĆ I ZRASZANIE	The skin dries out quickly, so the humidity over the land section must be 80-100 %. Mist the land section with dechlorinated water once or twice a day, and let moss and soil hold the moisture without the substrate becoming sodden.
WODA I JAKOŚĆ WODY	The water section is 10-20 cm deep and 10-20 °C. Use dechlorinated tap water or rainwater; the chlorine must be gone, but water quality is otherwise not critical.
OŚWIETLENIE I UVB	UVB is not necessary for the species, which lives hidden.
ŻYWIENIE	*Lissotriton boscai* eats any small animal it can swallow. In the water it is given daphnia, mosquito larvae, tubifex and small earthworms, and it also takes frozen food from the bottom.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is peaceful and can be kept in pairs or in a group with more females than males, so the females are left in peace in the breeding season.
NATURALNY ZASIĘG	Portugalia i Hiszpania.