herpy

SALAMANDRY I TRASZKI

Southern Banded Newt

Ommatotriton vittatus

Ommatotriton vittatus is a medium-sized newt of 9-11 cm from the Levant and southern Turkey. In the breeding season the male develops a tall, serrated dorsal crest and stronger colours. The species likes cool conditions, is shy, and needs a cool, humid and quiet set-up, so it is one of the more difficult newts to keep. The population in Turkey is declining, so captive-bred animals are the right choice.

Rodzina: Salamandridae Dawna nazwa: *Salamandra vittatus; Triturus vittatus* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**9-11 cm** długość całkowita z ogonem
Długość życia	**8-12 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Półwodny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**60 × 30 × 30 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**10-18 °C**
Temperatura w nocy	**5-12 °C**
Wilgotność	**80-100%**
Woda	**Akwaterrarium** głębokość 10-20 cm
Temperatura wody	**8-18 °C**
UVB	**Niewymagane**
Zimowanie	**4-10 °C** 8-12 tygodni
Pokarm	**Owady, małe bezkręgowce (skoczogonki, muszki owocowe) i dżdżownice i ślimaki**
Utrzymanie	**W parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**W wodzie**
Dojrzałość płciowa	**36-48 miesięcy**
Liczba jaj w zniesieniu	**100-200**
Okres larwalny	**60-120 dni** od wyklucia do metamorfozy
Pokarm larw	**Małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about keeping *Ommatotriton vittatus*.
TEMPERATURA	*Ommatotriton vittatus* likes cool conditions. The daytime temperature should be 10-18 °C and the night temperature 5-12 °C, and the temperature must not exceed 20 °C for long.
WILGOTNOŚĆ I ZRASZANIE	Keep the land section damp, with a humidity of 80-100 %, but never waterlogged. Mist the moss with dechlorinated water when it begins to dry.
WODA I JAKOŚĆ WODY	The water section should be 10-20 cm deep and 8-18 °C. Use dechlorinated tap water or rainwater. Clean water is important: change a third of the water weekly, and remove food remains daily. A small filter with a gentle flow helps.
OŚWIETLENIE I UVB	UVB is not necessary for the species, which is active at night and at dusk and avoids sun. Ordinary daylight or LED light for 10-12 hours a day is enough. The lamp must not heat the enclosure.
ŻYWIENIE	*Ommatotriton vittatus* eats any small animal it can overpower. On land, springtails, woodlice, white worms, fruit flies and small snails and spiders are suitable, and for adults also larger woodlice and crickets.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is shy and does best when left in peace. Males bite and chase each other away from their territory, so two males do not belong together in a small enclosure.
NATURALNY ZASIĘG	Turcja, Syria, Liban, Izrael i Jordania.