herpy

SALAMANDRY I TRASZKI

California Newt

Taricha torosa

Taricha torosa is a sturdy newt, 13-20 cm long, from the California coast, with a granular brown back and a yellow to orange belly. The skin contains tetrodotoxin, a nerve toxin with no antidote, and the animal can be deadly if swallowed. The species lives mostly on land and enters the water only to breed, and it must be kept cool.

Rodzina: Salamandridae Dawna nazwa: *Cynops torosus; Notophthalmus torosus* **Silne toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**13-20 cm** długość całkowita z ogonem
Długość życia	**12-22 lat**
Aktywność	**Za dnia i w nocy**
Tryb życia	**Półwodny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Silne** Unikaj kontaktu ze skórą i dokładnie umyj ręce po pracy w terrarium
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**60 × 30 × 30 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**16-21 °C**
Temperatura w nocy	**14-18 °C**
Wilgotność	**70-90%**
Woda	**Miska z wodą**
UVB	**Niewymagane**
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**W wodzie**
Dojrzałość płciowa	**36 miesięcy**
Liczba jaj w zniesieniu	**7-30**
Czas do wyklucia	**14-21 dni**
Okres larwalny	**60-90 dni** od wyklucia do metamorfozy
Pokarm larw	**Małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Bliski zagrożenia (NT)**

Hodowla w skrócie

TERRARIUM	There is relatively little knowledge about the care of *Taricha torosa*, so the winter temperatures are borrowed from the closely related *Taricha granulosa*, which is kept more often.
TEMPERATURA	*Taricha torosa* likes the cool. By day the temperature should be 16-21 °C and at night 14-18 °C. The temperature must not exceed 24 °C, and longer periods above 21 °C must be avoided.
WILGOTNOŚĆ I ZRASZANIE	The substrate should be damp, but the enclosure should not drip. Humidity is about 70-90%, and daily misting with dechlorinated water keeps the substrate moist.
WODA I JAKOŚĆ WODY	A shallow dish of clean water must always be available, so that the animal can cover itself completely with water. Use tap water treated for chlorine and chloramine; distilled and demineralised water are not used.
OŚWIETLENIE I UVB	The species does not need UVB. Daylight in the room or a weak fluorescent tube during the day and darkness at night is enough.
ŻYWIENIE	*Taricha torosa* is an opportunistic predator that takes almost anything it can overpower. Earthworms are a complete diet in themselves, and compost worms (*Eisenia fetida*) are also taken readily.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Taricha torosa* is calm, sociable and not aggressive towards others of its kind of the same size.
NATURALNY ZASIĘG	Stany Zjednoczone.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/salamanders/taricha-torosa