herpy

SALAMANDRY I TRASZKI

Danube Crested Newt

Triturus dobrogicus

Triturus dobrogicus is a slender crested newt, 13-18 cm long, from the Danube basin and the plains of the large rivers. It is the smallest and most aquatic of the crested newts, with an orange to red belly and sharply defined black spots. The male develops a serrated crest in spring. The species is rare in captivity, and its care follows that of the other crested newts.

Rodzina: Salamandridae

Słabe toksyny skórne

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**13-18 cm** długość całkowita z ogonem
Długość życia	**7-17 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Półwodny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**45 × 30 × 30 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**12-20 °C**
Temperatura w nocy	**8-15 °C**
Wilgotność	**70-95%**
Woda	**Akwaterrarium** głębokość 15-30 cm
Temperatura wody	**10-20 °C**
UVB	**Niewymagane**
Zimowanie	**4-10 °C** 6-12 tygodni
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**W wodzie**
Dojrzałość płciowa	**12-36 miesięcy**
Liczba jaj w zniesieniu	**200**
Czas do wyklucia	**14-35 dni**
Okres larwalny	**60-120 dni** od wyklucia do metamorfozy
Pokarm larw	**Gotowa karma, inne larwy i małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Triturus dobrogicus*, and the species is rare in captivity.
TEMPERATURA	*Triturus dobrogicus* loves cool conditions. The water is kept at 10-20 °C and must never become warmer; the animals are active down to about 5 °C.
WILGOTNOŚĆ I ZRASZANIE	In the aquatic phase humidity is not a problem. In the land phase the soft skin dries out quickly, so the substrate must be damp, with a humidity of 70-95%.
WODA I JAKOŚĆ WODY	Water depth is 15-30 cm, and the water is kept at 10-20 °C. The species is used to river water and tolerates many kinds of water, but chlorine harms it; tap water that has stood for a couple of days is unproblematic.
OŚWIETLENIE I UVB	The animals are nocturnal and do not need UVB. The plants need light to grow, and an ordinary aquarium lamp is enough.
ŻYWIENIE	In the water *T. dobrogicus* takes earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms. Frozen bloodworms and fish food are also taken, but often have to be offered with tweezers at first.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Triturus dobrogicus* is nocturnal and hides in vegetation by day. In the water the animals snap at anything that moves, including each other during feeding, and in the wild they also eat smaller newts and tadpoles.
NATURALNY ZASIĘG	Austria, Czechy, Słowacja, Węgry, Słowenia, Chorwacja, Bośnia i Hercegowina, Serbia, Rumunia, Ukraina, Bułgaria i Mołdawia.