herpy

SALAMANDRY I TRASZKI

Southern Crested Newt

Triturus karelinii

Triturus karelinii is a large, robust crested newt of 12-18 cm from the Crimea, the Caucasus and the southern shore of the Caspian Sea. It has brown to grey, warty skin, and the throat and belly are orange or yellow with small black spots. In spring the male develops a tall, serrated crest. The species is hardy and easy to keep, and it can be kept both on land and in water.

Rodzina: Salamandridae Dawna nazwa: *Triturus cristatus karelinii* **Słabe toksyny skórne**

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**12-18 cm** długość całkowita z ogonem
Długość życia	**8-11 lat**
Aktywność	**O zmierzchu i w nocy**
Tryb życia	**Półwodny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Toksyny skórne	**Słabe** Umyj ręce po kontakcie i trzymaj palce z dala od oczu i ust
Trudność	**Początkujący**

HODOWLA

Min. terrarium	**45 × 30 × 30 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**12-20 °C**
Temperatura w nocy	**8-15 °C**
Wilgotność	**70-95%**
Woda	**Akwaterrarium** głębokość 15-30 cm
Temperatura wody	**10-20 °C**
UVB	**Niewymagane**
Zimowanie	**4-10 °C** 6-12 tygodni
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Larwy w wodzie**
Miejsce składania jaj	**W wodzie**
Dojrzałość płciowa	**12-36 miesięcy**
Liczba jaj w zniesieniu	**150-400**
Czas do wyklucia	**14-35 dni**
Okres larwalny	**56-112 dni** od wyklucia do metamorfozy
Pokarm larw	**Gotowa karma, inne larwy i małe bezkręgowce wodne (rozwielitki, larwy komarów)**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	*Triturus karelinii* can be kept following its natural cycle, with an aquatic phase and a terrestrial phase, or in water all year.
TEMPERATURA	The water should be kept below 25 °C, because in warmer water the animals try to leave it. The animals are fully active down to about 10 °C.
WILGOTNOŚĆ I ZRASZANIE	In the aquatic phase humidity is not an issue. In the terrestrial phase the soft skin dries out quickly, so the substrate must be kept damp, with a humidity of 70-95%.
WODA I JAKOŚĆ WODY	A water depth of 15 cm is enough, and deeper is fine: 15-30 cm gives the animals room to swim. The water is kept at 10-20 °C.
OŚWIETLENIE I UVB	The animals are nocturnal and do not need UVB or artificial light for their own sake. The plants need light to grow, and an ordinary aquarium lamp is enough.
ŻYWIENIE	In the aquatic phase *T. karelinii* takes a wide range of food: earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Triturus karelinii* is robust and easy to keep. In the water the animals snap hard at anything that moves, including each other during feeding, and in the wild crested newts also eat smaller amphibians.
NATURALNY ZASIĘG	Ukraina, Rosja, Gruzja, Azerbejdżan i Iran.