herpy

SCYNKI

South-western Crevice Skink

Egernia napoleonis

A medium-sized, flat-bodied, diurnal skink from the rocks and forests of south-western Australia. *Egernia napoleonis* hides in crevices under bark and stones and gives birth to live young. Almost no documented knowledge exists on keeping it in captivity, so this guide draws on the species in the wild and on the close relatives *Egernia striolata*, *Egernia cunninghami* and *Egernia kingii*.

Rodzina: Scincidae Dawna nazwa: *Tiliqua napoleonis; Contundo napoleonis; Egernia bos*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**13-14 cm** długość od pyska do kloaki
Długość życia	**10-20 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny i skalny**
Środowisko	**Tereny skaliste i lasy i łąki strefy umiarkowanej**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**100 × 50 × 60 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**25-30 °C**
Miejsce wygrzewania	**35-40 °C**
Temperatura w nocy	**14-20 °C**
Wilgotność	**40-60%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady, owoce i rośliny**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Żyworodny**
Wielkość miotu	**1-6 młodych**
Zniesienia w roku	**1**
Ciąża	**90-120 dni**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik III**
Załącznik unijny	**Załącznik C**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Almost no documented knowledge exists on the care of *Egernia napoleonis*.
TEMPERATURA	By day the air should be 25-30 °C with a clear gradient from the cool end to the warm end. Under a halogen spot lies a flat stone or branch whose surface should reach 35-40 °C, measured with an infrared thermometer.
WILGOTNOŚĆ I WODA	Humidity should be 40-60 %. Mist the enclosure lightly a couple of times a week, and keep the substrate slightly damp under one hide in the cool end, so the animal can choose. A dish of fresh water should always be available.
OŚWIETLENIE I UVB	*Egernia napoleonis* is diurnal and needs UVB for as long as the light is on. A UVB tube in a reflector over the basking area, covering about half the length of the enclosure, gives Ferguson Zone 3.
ŻYWIENIE	The diet is not documented for *Egernia napoleonis*, so feeding follows the close relatives: an omnivorous diet with plenty of insects and a good deal of plant material.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Egernia napoleonis* is a diurnal skink that uses crevices as hides and climbs on bark and rocks. Close relatives in the same group, such as *Egernia striolata*, live in family groups.
NATURALNY ZASIĘG	Australia.