herpy

SCYNKI

Algerian orange-tailed skink

Eumeces algeriensis

A large, sturdy skink from Morocco and Algeria with a brown ground colour and pale crossbands with reddish eyespots. *Eumeces algeriensis* burrows into dry soil, needs sun, UVB and a deep substrate, and is often confused with *Eumeces schneiderii*.

Rodzina: Scincidae Dawna nazwa: *Eumeces schneideri algeriensis; Novoeumeces algeriensis*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**30-45 cm** długość całkowita z ogonem
Długość życia	**15-20 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny i kopiący nory**
Środowisko	**Półpustynia i lasy i łąki strefy umiarkowanej**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**120 × 60 × 45 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**24-30 °C**
Miejsce wygrzewania	**38-45 °C**
Temperatura w nocy	**21-24 °C**
Wilgotność	**20-40%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady, rośliny i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**5-15 jaj**
Inkubacja	**40-60 dni** przy 28-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM

Eumeces algeriensis lives on the ground and burrows, so floor area and substrate matter most. Two animals need at least 120 × 60 × 45 cm (length × width × height); a single animal can be kept in 90 × 45 × 40 cm.

TEMPERATURA

The enclosure needs a clear gradient. By day the temperature should be 24-30 °C, coolest at one end and warmest near the basking spot.

WILGOTNOŚĆ I WODA

The species comes from dry regions and must be kept dry: the humidity should stay below 40 %. Mist lightly once a day if the air is very dry, and twice a day when a shed is approaching and the skin looks dull.

OŚWIETLENIE I UVB

The species is diurnal and needs UVB for as long as the light is on. It is placed in Ferguson zone 3; the figure is borrowed from *Eumeces schneiderii*.

ŻYWIENIE

Eumeces algeriensis is omnivorous but feeds mainly on insects: crickets, cockroaches, mealworms, silkworms, waxworms and hornworms, as well as snails and earthworms.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

The species is less sociable than *E. schneiderii*. It can be kept singly, as a pair or as two females, but never two males together. The enclosure needs several hides and basking spots. New animals are often wary.

NATURALNY ZASIĘG

Maroko i Algieria.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/skinks/eumeces-algeriensis