herpy

SCYNKI

African fire skink

Mochlus fernandi

A heavy, fast skink from the forests of West Africa with vivid red-orange flanks, black and white crossbars and a bronze back. *Mochlus fernandi* burrows into moist substrate, needs warmth, UVB and moist substrate, and can live for 15-20 years.

Rodzina: Scincidae Dawna nazwa: *Lepidothyris fernandi; Lygosoma fernandi; Riopa fernandi; Tiliqua fernandi*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**25-38 cm** długość całkowita z ogonem
Waga	**50-100 g**
Długość życia	**15-20 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny i kopiący nory**
Środowisko	**Sawanna i las deszczowy**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**120 × 60 × 60 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**25-30 °C**
Miejsce wygrzewania	**33-36 °C**
Temperatura w nocy	**20-24 °C**
Wilgotność	**60-70%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 2
Pokarm	**Owady**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**3-9 jaj**
Zniesienia w roku	**3-5**
Inkubacja	**55-65 dni** przy 30 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM

Mochlus fernandi lives on the ground and burrows, so floor area and substrate matter most. A single animal can be kept in 90 × 45 × 45 cm, while a pair needs at least 120 × 60 × 60 cm (length × width × height).

TEMPERATURA

The enclosure needs a clear gradient. By day the temperature should be 25-30 °C, and at the basking spot the surface should reach 33-36 °C, measured with an infrared thermometer. The cool end should sit at about 24-26 °C.

WILGOTNOŚĆ I WODA

The species needs high humidity in the substrate: 60-70 % around the animal's hiding places, while the surface in the warm end may be drier, down to about 40 %.

OŚWIETLENIE I UVB

The species is diurnal and belongs to Ferguson zone 2: it basks a little and needs moderate UVB. The UV index should be about 3 at the basking spot and lower elsewhere in the enclosure.

ŻYWIENIE

Mochlus fernandi is an insectivore. The staple is crickets, supplemented with cockroaches, mealworms, silkworms, snails and other insects of a suitable size. Prey should be no larger than the animal's head.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

The species is fast, slippery and somewhat shy. A new animal needs at least two weeks of peace before it is handled, and it should be lifted gently from below in a closed room so that it cannot escape.

NATURALNY ZASIĘG

Sierra Leone, Gwinea, Liberia, Côte d'Ivoire, Ghana, Nigeria, Kamerun, Gwinea Równikowa, Gabon i Kongo.