herpy

WARANY

Large-scaled water monitor

Varanus nuchalis

Varanus nuchalis is a large, powerfully built water monitor from the central Visayan islands of the Philippines, formerly regarded as a subspecies of *Varanus salvator*. It is recognised by its greatly enlarged nuchal scales and is either marked with pale spots or almost black. Almost nothing is known about the care of this particular species, so much of the advice is borrowed from related water monitors.

Rodzina: Varanidae Dawna nazwa: *Varanus salvator nuchalis; Hydrosaurus nuchalis*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**100-150 cm** długość całkowita z ogonem
Długość życia	**15-20 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny, naziemny i półwodny**
Środowisko	**Las deszczowy i mokradła**
Trudność	**Ekspert**

HODOWLA

Min. terrarium	**300 × 150 × 150 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**27-32 °C**
Miejsce wygrzewania	**52-60 °C**
Temperatura w nocy	**21-24 °C**
Wilgotność	**60-70%**
Woda	**Część wodna** głębokość 35-50 cm
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady, kręgowce i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**5-10 jaj**
Zniesienia w roku	**1-3**
Inkubacja	**187-220 dni** przy 27-31,5 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM

Little is known about the care of *Varanus nuchalis*. The figures are borrowed from close relatives in the same group, especially *Varanus cumingi* and *Varanus salvator*.

TEMPERATURA

Varanus nuchalis needs a background temperature of 27-32 °C by day. The basking spot should have a surface temperature of 52-60 °C, measured with an IR thermometer on the basking spot itself, not in the air.

WILGOTNOŚĆ I WODA

Humidity should be 60-70 %. A large water basin keeps humidity up on its own, and a damp, deep substrate gives the animal somewhere to retreat to.

WODA I JAKOŚĆ WODY

The species is tied to water, so a large water section is not optional. It must be big enough for the whole animal to lie submerged and swim, and the depth should be 35-50 cm for an adult.

OŚWIETLENIE I UVB

The species is diurnal and basks, so strong light and UVB are essential. It belongs to Ferguson zone 3, like other diurnal monitors that bask a great deal.

ŻYWIENIE

The diet of *Varanus nuchalis* in the wild has not been studied. Related water monitors eat many kinds of animals and carrion, and the diet below is borrowed from them.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

Varanus nuchalis must be handled like other large water monitors. Even a medium-sized monitor has sharp teeth, strong claws and a powerful tail that it can lash with. A bite can cause deep wounds, and the claws can tear the skin.

NATURALNY ZASIĘG

Filipiny.