herpy

WĘŻE

Plains Garter Snake

Thamnophis radix

Thamnophis radix is a slim, striped garter snake from the prairies and open grasslands of central North America. It has a yellow or orange stripe down the back and grows to 40-70 cm. It is very tolerant of cold and gives birth to live young. It is hardy and can be kept on thawed mice, but it escapes through the smallest gaps.

Rodzina: Colubridae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**40-70 cm** długość całkowita z ogonem
Długość życia	**5-10 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Mokradła i lasy i łąki strefy umiarkowanej**
Trudność	**Początkujący**

HODOWLA

Min. terrarium	**100 × 50 × 50 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**23-27 °C**
Miejsce wygrzewania	**29-32 °C**
Temperatura w nocy	**18-24 °C**
Wilgotność	**35-60%**
Woda	**Miska z wodą**
UVB	**Zalecane** Strefa Fergusona 2
Zimowanie	**8-15 °C** 8-16 tygodni
Pokarm	**Kręgowce i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Żyworodny**
Dojrzałość płciowa	**24-36 miesięcy**
Wielkość miotu	**5-60 młodych**
Zniesienia w roku	**1**
Ciąża	**83-102 dni**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little species-specific knowledge exists about keeping *Thamnophis radix*.
TEMPERATURA	There must be a clear temperature gradient. The basking spot should be 29-32 °C measured on the surface, the cool end 23-27 °C and the night 18-24 °C.
WILGOTNOŚĆ I WODA	The air humidity should be 35-60 %, highest at the cool end and at night. The enclosure should be dry, because persistently damp conditions cause blisters on the belly and fungal infections.
OŚWIETLENIE I UVB	*Thamnophis radix* is diurnal and thrives in strong daylight. UVB is not strictly necessary, but is recommended.
ŻYWIENIE	In the wild *Thamnophis radix* eats earthworms, slugs, frogs, toads, salamanders, fish and the occasional mouse.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Thamnophis radix* is quick and alert, but usually becomes calm if handled regularly from a young age. Let new animals settle in for about two weeks, and start handling them only when they are feeding steadily.
NATURALNY ZASIĘG	Kanada i Stany Zjednoczone.