herpy

ŻABY

Haraldmeier's Mantella

Mantella haraldmeieri

Mantella haraldmeieri is a small, diurnal frog of 21-27 mm from the south-eastern forests of Madagascar, light brown on the back with black flanks, copper-coloured hind legs and red markings on the underside of the legs. The species is threatened in the wild and only rarely kept, so very little is known about its care. It is kept like the other species in the genus: cool, humid and never warm.

Rodzina: Mantellidae Dawna nazwa: *Mantella madagascariensis haraldmeieri*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**21-27 mm** długość od pyska do kloaki
Długość życia	**5-10 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Las deszczowy**
Toksyny skórne na wolności	**Słabe**
Toksyny skórne w niewoli	**Brak** Zwierzęta z hodowli. Osobniki odłowione na wolności mogą przez długi czas zachować toksyny
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**45 × 45 × 30 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**21-24 °C**
Temperatura w nocy	**16-18 °C**
Wilgotność	**60-80%**
Woda	**Miska z wodą**
UVB	**Opcjonalne** Strefa Fergusona 1
Pokarm	**Małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**W grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Kijanki w wodzie**
Miejsce składania jaj	**Na ziemi**
Dojrzałość płciowa	**12-18 miesięcy**
Liczba jaj w zniesieniu	**50-130**
Czas do wyklucia	**3-12 dni**
Okres larwalny	**42-56 dni** od wyklucia do metamorfozy
Pokarm larw	**Glony i rośliny i gotowa karma**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
IUCN	**Zagrożony (EN)**

Hodowla w skrócie

TERRARIUM	Very little is known about keeping *Mantella haraldmeieri*, because the species is only rarely kept.
TEMPERATURA	The temperature should be 21-24 °C during the day and 16-18 °C at night, corresponding to a cool room. Heat should be avoided: over 27 °C can quickly become fatal for the genus. The figures are borrowed from *M. baroni*.
WILGOTNOŚĆ I ZRASZANIE	Humidity should be 60-70 % day to day and up to 80 % in the breeding season (borrowed from *M. baroni*). Mist the enclosure daily with decalcified water, heavily in warm weather and a little less when it is cool.
WODA I JAKOŚĆ WODY	Place a shallow water dish in the enclosure, and keep it full at all times. The frogs are poor swimmers, so the dish must be so shallow that they can easily get out, and it must not take up much of the floor.
OŚWIETLENIE I UVB	No UVB requirement is documented for the species. The genus is active in partial sunlight in the wild, so a weak UVB light (Ferguson zone 1) is probably an advantage, provided there is shade and plants so the animals can choose.
ŻYWIENIE	*Mantella haraldmeieri* is fed small arthropods: springtails, wingless fruit flies, newly hatched crickets, small woodlice, aphids and lesser mealworm larvae. The genus takes prey of up to about 6 mm. Variety prevents deficiencies.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The frog is diurnal and patrols its territory when hunting. The males call, defend territories and may wrestle with one another.
NATURALNY ZASIĘG	Madagaskar.