herpy

ŻABY

Marojejy Mantella

Mantella manery

Mantella manery is a small, diurnal frog from the forests of northern Madagascar, dark brown with a white stripe on the side of the head. It is a close relative of *Mantella laevigata*. The species is vulnerable in the wild and known from few records, and little is known about its care, so the figures here are borrowed from related species.

Rodzina: Mantellidae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**2-3 cm** długość od pyska do kloaki
Długość życia	**4-8 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Las deszczowy i mokradła**
Toksyny skórne na wolności	**Słabe**
Toksyny skórne w niewoli	**Brak** Zwierzęta z hodowli. Osobniki odłowione na wolności mogą przez długi czas zachować toksyny
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**45 × 45 × 30 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**20-26 °C**
Temperatura w nocy	**17-22 °C**
Wilgotność	**80-100%**
Woda	**Miska z wodą**
UVB	**Opcjonalne** Strefa Fergusona 1
Pokarm	**Małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**W grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Kijanki w wodzie**
Miejsce składania jaj	**Na ziemi**
Dojrzałość płciowa	**12-18 miesięcy**
Liczba jaj w zniesieniu	**12-30**
Czas do wyklucia	**3-12 dni**
Okres larwalny	**42-98 dni** od wyklucia do metamorfozy
Pokarm larw	**Glony i rośliny i gotowa karma**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
IUCN	**Narażony (VU)**

Hodowla w skrócie

TERRARIUM	Little is known about keeping *Mantella manery*. The habitat is leaf litter by water in low-lying forest in northern Madagascar.
TEMPERATURA	The temperature should be 20-26 °C during the day and 17-22 °C at night. The figures are borrowed from *M. laevigata*, which also comes from the lowlands and tolerates somewhat more heat than most species in the genus.
WILGOTNOŚĆ I ZRASZANIE	Humidity should be 80-100 % (borrowed from *M. laevigata*). Mist the enclosure daily with decalcified water, heavily in warm weather and a little less when it is cool. The substrate should be damp, but not waterlogged.
WODA I JAKOŚĆ WODY	Place a shallow water dish in the enclosure, and keep it full at all times. The frogs are poor swimmers, so the dish must be so shallow that they can easily get out, and it must not take up much of the floor.
OŚWIETLENIE I UVB	The UVB requirement is not documented for the species. A weak UVB light in Ferguson zone 1 is an advantage when there is shade and plants so the animals can choose for themselves. Light for the plants should not give off heat.
ŻYWIENIE	The genus takes prey of up to about 6 mm. Give newly hatched crickets, wingless fruit flies, springtails, small woodlice, aphids, small cockroach nymphs, lesser mealworm larvae and termites.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The frogs of the genus are diurnal. The males call to defend territories and attract females, and they may wrestle with one another. The species' own behaviour is not documented.
NATURALNY ZASIĘG	Madagaskar.