herpy

ŻABY

Chantaburi Bug-eyed Frog

Theloderma stellatum

Theloderma stellatum is a small, tree-dwelling mossy frog from the lowland rainforest of Thailand, Cambodia, Laos and Vietnam. It resembles a darker version of *Theloderma asperum*, grows to about 3-4 cm and lays its eggs above water in water-filled tree hollows. Its care is similar to that of its relatives, but almost all figures are borrowed from them, because the species is rare in captivity.

Rodzina: Rhacophoridae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**3-4 cm** długość od pyska do kloaki
Długość życia	**5-10 lat**
Aktywność	**W nocy**
Tryb życia	**Nadrzewny i półwodny**
Środowisko	**Las deszczowy**
Toksyny skórne	**Brak znanych**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**45 × 45 × 45 cm** D × S × W dla 4 dorosłych osobników
Temperatura w dzień	**20-24 °C**
Temperatura w nocy	**17-21 °C**
Wilgotność	**75-95%**
Woda	**Część wodna** głębokość 5-10 cm
UVB	**Opcjonalne** Strefa Fergusona 1
Pokarm	**Owady**
Utrzymanie	**W parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Kijanki w wodzie**
Miejsce składania jaj	**Na liściach nad wodą**
Dojrzałość płciowa	**8-10 miesięcy**
Liczba jaj w zniesieniu	**4-15**
Czas do wyklucia	**10-14 dni**
Okres larwalny	**75-180 dni** od wyklucia do metamorfozy
Pokarm larw	**Gotowa karma**
Opieka nad potomstwem	**Brak**

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little documented knowledge exists on the care of *Theloderma stellatum*.
TEMPERATURA	By day the temperature should be 20-24 °C and at night 17-21 °C.
WILGOTNOŚĆ I ZRASZANIE	Humidity should be 75-95 % and never below 60 %, otherwise the frogs may dry out. The large water surface keeps the humidity up on its own, so the frogs do not need much misting.
WODA I JAKOŚĆ WODY	The frog spends a lot of time at the water's edge and even sits under water, so the water section is not there for show.
OŚWIETLENIE I UVB	The frog is nocturnal and has no documented need for UVB. A weak UVB tube for shade species (Ferguson zone 1) can be used and is optional, but the frogs must always be able to retreat into shade.
ŻYWIENIE	*T. stellatum* eats insects. Small crickets are the main food, supplemented with waxworms, flies, springtails for the smallest young, small dubia roaches and soldier fly larvae.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Theloderma stellatum* is nocturnal and sits hidden by day in tubes, among plants or close to the water. The frog does well in a group as long as there is enough space. The males call at night.
NATURALNY ZASIĘG	Tajlandia, Kambodża, Laos i Wietnam.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/frogs/theloderma-stellatum