herpy

ŻÓŁWIE LĄDOWE I WODNE

South African Bowsprit Tortoise

Chersina angulata

Chersina angulata is a medium-sized, domed tortoise from the coastal and scrub regions of South Africa. It is recognised by the large, projecting shield under the chin (the gular scute), which males use to overturn rivals. The shell length is typically 17-19 cm. It is hardy in its own Mediterranean-type climate, but copes poorly with cold and damp and needs plenty of space, sun and dry, sandy ground.

Rodzina: Testudinidae Dawna nazwa: *Testudo angulata; Testudo bellii; Goniochersus angulatus; Neotestudo angulata; Chersine angulata*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**15-35 cm** długość całkowita z ogonem
Długość życia	**30-50 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Półpustynia**
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**160 × 80 × 40 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**22-28 °C**
Miejsce wygrzewania	**32-35 °C**
Temperatura w nocy	**12-18 °C**
Wilgotność	**40-60%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady i rośliny**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Dojrzałość płciowa	**120-144 miesięcy**
Wielkość zniesienia	**1-2 jaj**
Zniesienia w roku	**1-6**
Inkubacja	**94-198 dni** przy 28-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little published knowledge exists on the care of *Chersina angulata* outside South Africa.
TEMPERATURA	By day the enclosure should be 22-28 °C, with a basking spot of 32-35 °C under a lamp, measured on the substrate. The figures are estimated from the climate of the habitat and borrowed from *T. hermanni*.
WILGOTNOŚĆ I WODA	*C. angulata* belongs to dry to semi-dry habitats and needs a humidity of 40-60 %; the figure is an estimate based on the habitat. Keep most of the substrate dry and make a slightly damper corner under a shelter.
OŚWIETLENIE I UVB	*C. angulata* is diurnal and basks a great deal. Sunlight is best, and the animal should be outside in the sun whenever the weather allows.
ŻYWIENIE	In the wild herbs and grasses are the most important food, and the animal also eats moss, fungi, insects, snail shells and droppings. It feeds selectively and changes its food with the season.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is shy and withdraws under bushes or half-buries itself.
NATURALNY ZASIĘG	Republika Południowej Afryki i Namibia.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/tortoises-and-turtles/chersina-angulata