herpy

ŻÓŁWIE LĄDOWE I WODNE

Forsten's Tortoise

Indotestudo forstenii

Indotestudo forstenii is a medium-sized tortoise from the forests of Sulawesi and Halmahera in Indonesia. The carapace is caramel to dark brown with black blotches, and adults typically reach about 25 cm. The species is crepuscular and shy, needs warmth and high humidity, and does not hibernate. It is critically endangered, and animals taken from the wild are often sick.

Rodzina: Testudinidae Dawna nazwa: *Testudo forstenii; Geochelone forsteni; Indotestudo elongata forsteni*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**25-30 cm** długość całkowita z ogonem
Waga	**2 000-3 000 g**
Długość życia	**30-50 lat**
Aktywność	**O zmierzchu**
Tryb życia	**Naziemny**
Środowisko	**Las deszczowy**
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**180 × 120 × 40 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**26-30 °C**
Miejsce wygrzewania	**30-32 °C**
Temperatura w nocy	**18-20 °C**
Wilgotność	**70-85%**
Woda	**Miska z wodą**
UVB	**Zalecane** Strefa Fergusona 2
Pokarm	**Owoce, rośliny i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**1-4 jaj**
Zniesienia w roku	**1-3**
Inkubacja	**130-165 dni** przy 26-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Krytycznie zagrożony (CR)**

Hodowla w skrócie

TERRARIUM

Little is known about the care of *Indotestudo forstenii*, and much is borrowed from the close relative *Indotestudo elongata*. The species needs a large, open floor area with damp substrate and hides.

TEMPERATURA

The air should be 26-30 °C by day and 18-20 °C at night. The animals do not seek the sun and seldom sit under the lamp, but a warm zone of 30-32 °C under a heat lamp at one end, measured at shell height, gives them a choice.

WILGOTNOŚĆ I WODA

Humidity should be high, about 70-85 %, with a damper and a drier zone so that the animal can choose. The substrate should be damp but not wet, and the hides are dampened regularly.

OŚWIETLENIE I UVB

Indotestudo forstenii avoids bright light, but needs vitamin D3 for calcium metabolism. The best solution is time outdoors in partial shade; glass and acrylic block UVB.

ŻYWIENIE

Indotestudo forstenii is omnivorous, with plants as the main component. The basis is leaves and herbs such as romaine lettuce, endive, dandelion and plantain together with grated vegetables.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

Indotestudo forstenii is shy and most active at dawn and dusk. It needs several hides and must not be disturbed all the time. Lift it only when necessary, under the plastron with both hands.

NATURALNY ZASIĘG

Indonezja.