herpy

ŻÓŁWIE LĄDOWE I WODNE

Common Spider Tortoise

Pyxis arachnoides

Pyxis arachnoides is a small, critically endangered tortoise from the dry spiny forests along the south-west coast of Madagascar. The shell is high and domed with a web-like pattern, and adults are 10-12 cm long and weigh 200-400 g. The species is rare in captivity and reproduces slowly, and little is known about its care.

Rodzina: Testudinidae Dawna nazwa: *Pyxis aranoides; Bellemys arachnoides*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**10-15 cm** długość całkowita z ogonem
Waga	**200-400 g**
Długość życia	**50-70 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Półpustynia i suchy las**
Trudność	**Ekspert**

HODOWLA

Min. terrarium	**120 × 60 × 40 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**24-30 °C**
Miejsce wygrzewania	**32-38 °C**
Temperatura w nocy	**18-22 °C**
Wilgotność	**50-70%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady i rośliny**
Utrzymanie	**Pojedynczo, w parach lub w grupach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**1 jaj**
Zniesienia w roku	**1-2**
Inkubacja	**220-330 dni** przy 29-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik I**
Załącznik unijny	**Załącznik A**
IUCN	**Krytycznie zagrożony (CR)**

Hodowla w skrócie

TERRARIUM	Little documented information exists on the care of *Pyxis arachnoides*. The species is kept and bred mainly in zoos and by specialists.
TEMPERATURA	The animal must be kept warm. During the day the temperature should be 24-30 °C, with a basking spot of 32-38 °C under a heat lamp at one end, measured on the substrate. At night the temperature drops to 18-22 °C.
WILGOTNOŚĆ I WODA	Humidity should be around 50-70 %. The figure is borrowed from *Astrochelys radiata* and should be seen as a starting point. Keep the substrate slightly damp, and mist the enclosure now and then.
OŚWIETLENIE I UVB	*P. arachnoides* is diurnal. Natural sunlight is best, and indoors the species needs UVB light with a basking spot in Ferguson zone 3. The zone is borrowed from *Astrochelys radiata* and has not been measured for the species.
ŻYWIENIE	In the wild *P. arachnoides* eats young leaves, flowers, fruit, insect larvae and the droppings of larger animals.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	Handle the animal as little as possible, especially young animals. When threatened it withdraws its head, legs and tail into the shell.
NATURALNY ZASIĘG	Madagaskar.