herpy

AGAMÍDEOS

Sailfin Lizard

Hydrosaurus amboinensis

Hydrosaurus amboinensis is a large, semi-aquatic agamid from the Moluccas and New Guinea. The male has a crest on the nape and a tall sail on the tail. It reaches 60-120 cm, basks on branches over the water and plunges into it when in danger. It needs a very large, warm and humid enclosure with a deep water section and powerful UVB.

Família: Agamidae Nome anterior: *Lacerta amboinensis; Lophura amboinensis*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**60-120 cm** comprimento total com a cauda
Longevidade	**15-25 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e semiaquático**
Habitat	**Floresta tropical e área úmida**
Dificuldade	**Especialista**

CUIDADOS

Terrário mín.	**200 × 80 × 150 cm** C × L × A para 1 adulto
Temperatura, dia	**26-32 °C**
Ponto quente	**35-45 °C**
Temperatura, noite	**24-27 °C**
Umidade	**75-90%**
Água	**Aquaterrário** profundidade 30-50 cm
UVB	**Essencial** Zona de Ferguson 3
Alimentação	**Insetos, frutas, plantas e vertebrados**
Mantido	**Sozinho, em casal ou um macho com várias fêmeas**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**2-8 ovos**
Incubação	**60-75 dias** a 28-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Sobre a espécie

Hydrosaurus amboinensis is a powerful, laterally flattened agamid with a tall body, a long tail and long toes. The male has a crest on the nape and a tall, upright sail on the front part of the tail, which is used for display and as a rudder and fin in the water. The colours are dark, olive and brown with lighter markings. The females are about a third smaller and have only a low crest and a low sail. The species was described in 1768 from animals from Ambon, which also gave it its name.

Adults typically reach 60-90 cm including the tail, and large males reach 105-120 cm. Together with *H. pustulatus*, the species is among the largest in the genus.

The genus *Hydrosaurus* comprises five species. Until 2020, the animals from Sulawesi were included in *H. amboinensis*, but then *H. microlophus* and *H. celebensis* were recognised as separate species, and *H. amboinensis* is now restricted to the central Moluccas and New Guinea. *H. weberi* from Halmahera has long been a separate species. Older literature and trade names

therefore use *H. amboinensis* for animals that today belong to other species, so have the origin stated when buying. There is almost no care information specific to the species; the care section is based on experience with the genus, especially *H. pustulatus*.

It is a nervous, fast and powerful lizard that is not suited to handling. It needs a very large enclosure, high heat, high humidity, UVB and a large water section, and it can live 15-25 years in captivity. The species is only for experienced keepers with plenty of space.

Saiba mais: Luz e UVB

Distribuição e habitat

Hydrosaurus amboinensis lives in the central Moluccas in eastern Indonesia, including Ambon and Seram, and on New Guinea, in both the Indonesian part and Papua New Guinea. The type locality is Ambon. The populations on Sulawesi now belong to *H. microlophus* and *H. celebensis*, and *H. weberi* lives on Halmahera, so older maps showing the species there are outdated. The exact distribution on New Guinea is poorly known.

> **Área de ocorrência natural:** Indonésia e Papua-Nova Guiné.

HABITAT

Hydrosaurus amboinensis lives in the lowlands along rivers and streams in tropical forest, in coastal forest and in mangroves. It is active by day and basks on branches that hang out over the water. When in danger, it runs down into the water, swims away and can stay on the bottom for a long time until the danger has passed. It also climbs in trees and shrubs by the water.

The climate is warm and humid all year round. The female buries her eggs in the soil near the water. The species is omnivorous and eats shoots, leaves, flowers, fruit, insects, crustaceans and small animals. The species' way of life in the wild has been little studied and resembles that of *H. pustulatus*.

Cuidados

O TERRÁRIO

An adult needs an enclosure of at least 200 × 80 × 150 cm (length × width × height). The dimensions are large because the animal needs a water section covering up to two thirds of the floor, a land section and high basking spots, and because an adult easily reaches a metre in length. This requires a home-built enclosure or a dedicated room; the standard enclosures sold in shops are too small. A pair or a trio needs each dimension to be at least 60 cm larger. Young animals can live in around 120 × 50 × 70 cm with a smaller water section, but quickly outgrow it.

The species lives both in the trees and in the water and often jumps into the water from a great height. Therefore furnish with horizontal and slanted branches that are at least as thick as the animal's body, and with shelves along the walls at different heights. The branches must hang out over the water, so that the animal can flee down into it. The water must be at least 30 cm deep, so that it breaks the fall; in shallow water the animal can injure itself. Make room on the land section for a food bowl and, if there are females, a laying site.

The substrate must tolerate high humidity without rotting: pure cypress bark, other bark mixes for moisture-loving reptiles or equal parts unfertilised soil, sand and moistened coconut fibre in a layer of a few centimetres. Cover all sides except the front, and fit a 30 cm high screen at the bottom of the front. This gives security and prevents the animal from rubbing its snout raw against the glass. The enclosure must be lockable, because the animal is fast and nervous.

TEMPERATURA

Hydrosaurus amboinensis is a tropical species and needs a clear gradient. The air is kept at 26-32 °C by day, and the basking spot above a favourite branch or a high shelf should have a surface temperature of 35-45 °C, measured with an infrared thermometer. Start at the low end and watch the animal: an animal that constantly seeks away from the basking spot or down into the water is too warm. At night the temperature falls to 24-27 °C.

Use halogen spotlights in ceramic fittings over the basking spot in the upper part of the enclosure, and put guards around them so that the animal cannot touch the bulb. Ceramic heaters retain heat at night and in the cold season without giving light. Heating cables under the substrate can help, but only if fitted according to the manufacturer's instructions, because

burns and fire are a real risk. Use a thermostat on all heat sources, and measure with digital thermometers both at the basking spot and in the cool part.

The animal must always be able to move away from the heat, and it readily cools off in the water.

UMIDADE E ÁGUA

The humidity should be at least 75 %, preferably 75-90 %. High humidity is essential for clean sheds, including on the crest, sail, spines and toes. Mist the enclosure at least once a day, or use a fogger or rain system, and keep the substrate evenly moist. The large water section evaporates constantly and helps a great deal.

Moisture and warmth make mould and bacteria grow. Provide ventilation both high and low, and replace the substrate when it turns sour or smells. Air that is too dry leaves old skin stuck on the crest, toes and tail.

ÁGUA E QUALIDADE DA ÁGUA

Hydrosaurus amboinensis uses the water for swimming, fleeing, cooling off and drinking. The water section must be large enough for the animal to dive and swim freely, and at least 30 cm deep; up to two thirds of the floor can be water. Make sure the animal can easily get out again by way of rocks, roots or a slope.

The animal defecates in the water and drinks from it, so the water must be changed daily, and the water section cleaned thoroughly at regular intervals. A powerful filter relieves the work, but does not replace the water change. Use clean tap water at room temperature.

ILUMINAÇÃO E UVB

Hydrosaurus amboinensis is active by day and basks, so it needs strong light and UVB. Use a powerful UVB tube (T5 HO, 10-12 %) in a reflective fitting covering a large part of the basking area. Ferguson zone 3 suits a diurnal species that basks at forest edges and along streams; the zone has not been established for the genus and is an estimate. Aim for a UV index of about 3-5 at the animal's back at the basking spot and lower in the shade, check with a UV meter, and replace the tube according to the manufacturer's instructions, typically every 12 months.

Mesh and glass dampen UVB, so mount the tube inside the enclosure with a guard around it, or over a coarse mesh. Provide 12 hours of light a day all year round, controlled by a timer. An extra daylight tube or an LED of 5,000-6,500 K gives a bright, natural enclosure.

ALIMENTAÇÃO

Hydrosaurus amboinensis is omnivorous. In the wild it eats shoots, leaves, flowers, fruit, insects, crustaceans and small vertebrates. In the enclosure, the diet consists of plants and a moderate amount of animal food.

Give plants every day, for example dandelion, stinging nettle, clover, rocket, kale, Chinese cabbage, herbs and edible flowers, supplemented with grated carrot, peas, beans and pumpkin and a little fruit such as berries, melon and banana. Chop it finely for juveniles, and leave the bowl out all day. Complete food for bearded dragons is a good addition.

Juveniles get insects (crickets, cockroaches, locusts, morio worms) every day or every other day and plants every day. Adults get insects a couple of times a week and occasionally a small mouse, prawns, freshwater crayfish or a cleaned fish. The insects should be well fed and dusted with calcium; give calcium with D3 and vitamins once or twice a week, and extra calcium for females that lay eggs. Too much animal food leads to obesity. The animal readily takes food from tongs, and that builds trust.

Saiba mais: Cálcio, vitaminas e suplementos

COMPORTAMENTO E MANUSEIO

Hydrosaurus amboinensis is nervous and jumpy. When in danger, it runs, often on its hind legs, down into the water and stays on the bottom until the danger has passed. In the enclosure, a frightened lizard throws itself at the glass and wears its snout. Screening, many hiding places and quiet around the enclosure are therefore the most important things.

The animal calms down with time, but it is not an animal for daily handling. Move slowly near the enclosure, and use hand feeding to build trust. If the animal has to be moved, lift it calmly from below instead of chasing it around: a chased animal is stressed, and the long claws scratch. Once it has been lifted, it normally settles quickly. It twists and scratches, but rarely bites.

Two males cannot be kept together. A pair or a male with two females can be kept in a very large set-up, but the animals must be able to be separated if stress or injuries arise. Like all reptiles, the animal can carry salmonella, so wash your hands after contact with the animal, the enclosure and the water.

Saiba mais: Quarentena e higiene

Reprodução

MATURIDADE SEXUAL E SEXAGEM

The males of *Hydrosaurus amboinensis* have a larger head, a taller crest and a much taller tail sail than the females, and they are about a third larger. In the female, the sail is only a low fringe. Young animals cannot be sexed reliably; the sexes only become clearly visible when the animals are well over half grown.

The age at sexual maturity is not documented. Only breed fully grown animals in good condition; a female that is too young or thin is worn down by egg laying.

ACASALAMENTO

Little is known about mating in *Hydrosaurus amboinensis*. The male displays with crest and sail and takes on stronger colours in the breeding season. In the wild, the young hatch in the rainy season. What triggers mating in captivity is not documented.

A pair or a male with two females requires a very large enclosure with many hiding places so that the females can withdraw from the male. Watch the animals, and separate them if a female gets wounds or stops eating. Give the females extra calcium and plenty of food before and after mating.

POSTURA E NINHADAS

The female lays 2-8 eggs per clutch; the figure is known from *H. pustulatus* and *H. weberi*, not from the species itself. She normally lays one clutch per season, in good years several. In the wild, she digs a shallow pit in the soil near the water, but above the flood line.

Provide the female with a laying box on the land section with moist, loose soil and sand, at least 30 cm deep, and cover it partly so that she feels secure. Without a suitable site she can retain the eggs, and that can become life-threatening; contact a vet if she seems strained or does not manage to lay. Females can lay infertile eggs without a male.

INCUBAÇÃO

Lift the eggs out carefully without turning them, and place them half buried in damp vermiculite or perlite (roughly equal parts material and water by weight) in a closed box. Incubate the eggs at a stable 28-30 °C. They hatch after 60-75 days; the figure is an estimate based on *H. weberi*, where the incubation period is around 65 days.

Check the box regularly for mould, and remove eggs that go mouldy or collapse.

CRIAÇÃO DOS FILHOTES

The hatchlings are independent from the first day and can run across the surface of the water on their flat toes. Put them in an enclosure of about 120 × 50 × 70 cm with branches, a small water section and many hiding places, and screen the sides so that they do not wear their snouts. Without a substrate the enclosure is easy to keep clean; the humidity is maintained with misting and water.

Feed every day with small insects, dusted with calcium, and finely chopped plants, and provide UVB from day one. The hatchlings grow fast and must be moved to larger enclosures as they grow. Males become territorial, so divide the young when fighting arises.

Saúde e erros comuns

Most problems in *Hydrosaurus amboinensis* are caused by the husbandry: too little heat, too little UVB, air that is too dry or a one-sided diet. Typical problems are:

- Brittle bones (metabolic bone disease) with too little UVB and calcium.
- Snout injuries when the animal runs into the glass. Screen the sides, and give peace.
- Skin stuck on the crest, toes and tail when the air is too dry.
- Burns from heat lamps without guards and from heating cables.
- Injuries after jumping into water that is too shallow.
- Skin and mouth infections with dirty water or a wet, sour substrate.
- Obesity with too much animal food.

- Egg-laying problems in females without a suitable laying site.
- Intestinal worms and other parasites, especially in wild-caught animals. Quarantine new animals, and have a faecal sample examined.

A healthy animal has clear eyes, a full body and tail and an intact crest and sail. It eats regularly, basks and seeks the water when frightened. Reptiles hide illness for a long time, so contact a vet experienced with reptiles if the animal loses weight, does not eat, has wounds or swellings, or sits with its mouth open.

Pergunte sempre a quem entende do assunto

Este guia é um bom lugar para começar, mas não substitui a experiência. Os animais diferem, e boa parte do que mais importa se aprende com quem cria a espécie há anos. Converse com um criador, um tutor experiente ou um veterinário que conheça répteis e anfíbios antes de ter o animal - e de novo se algo parecer errado.