herpy

ANGUÍDEOS

Fringed Arboreal Alligator Lizard

Abronia fimbriata

Abronia fimbriata is a brown, arboreal alligator lizard with thorn-like scales above the ear, found in the cloud forests of Guatemala's north-eastern highlands. It reaches about 25-35 cm, gives birth to live young and eats insects. It needs a tall, airy enclosure with moss and bromeliads and cool temperatures. The species is threatened, and its care is only sparsely documented.

Família: Anguidae Nome anterior: *Barissia fimbriata; Gerrhonotus fimbriatus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**25-35 cm** comprimento total com a cauda
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola**
Habitat	**Floresta de montanha**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**50 × 50 × 75 cm** C × L × A para 2 adultos
Temperatura, dia	**20-27 °C**
Ponto quente	**26-30 °C**
Temperatura, noite	**10-18 °C**
Umidade	**60-80%**
Água	**Sem recipiente de água**
UVB	**Recomendado** Zona de Ferguson 2
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Vivíparo**
Maturidade sexual	**24-36 meses**
Tamanho da ninhada	**4 filhotes**
Posturas por ano	**1**
Gestação	**180-240 dias**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice I**
Anexo da UE	**Anexo A**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	Care of *Abronia fimbriata* is only sparsely documented, and the figures given here are borrowed from the closely related *Abronia graminea*, which is the best-known species in the genus.
TEMPERATURA	*Abronia fimbriata* does not tolerate heat. The daytime temperature should be 20-27 °C, and the basking spot under a weak spotlight may reach 26-30 °C. The animal is most active at 24-27 °C.
UMIDADE E ÁGUA	Relative humidity should be 60-80 %, with the highest values at night and after misting. Mist the plants, walls and animals lightly almost every day, ideally with a pressure sprayer or a fogger that produces a fine mist.
ILUMINAÇÃO E UVB	The species is diurnal and needs UVB. The best option is natural, unfiltered sunlight, for example in an outdoor mesh cage in dry weather in summer, with shade and a cool surface so that the animal can move away.
ALIMENTAÇÃO	*Abronia fimbriata* feeds on insects and other small animals. Offer crickets, grasshoppers, mealworms, cockroaches, snails, spiders and soft larvae, for example green caterpillars and fly larvae with a high calcium content.
COMPORTAMENTO E MANUSEIO	The species is quiet and shy. The animal spends most of the day sitting still on a branch or in a bromeliad and basks for short periods. It is not an animal to be handled.
ÁREA DE OCORRÊNCIA NATURAL	Guatemala.