herpy

CAMALEÕES

Böhme's Two-horned Chameleon

Kinyongia boehmei

Kinyongia boehmei is a small, green mountain chameleon found only in the Taita Hills of south-eastern Kenya. The male has two flat, almost parallel nose horns, while the female is hornless. The species lays eggs and needs a cool, humid climate and a tall, densely planted enclosure. Its care is poorly documented, and it suits the experienced keeper best.

Família: Chamaeleonidae Nome anterior: *Bradypodion tavetanum boehmei; Kinyongia tavetana boehmei*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**15-19 cm** comprimento total com a cauda
Longevidade	**6-8 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola**
Habitat	**Floresta de montanha**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**45 × 45 × 90 cm** C × L × A para 1 adulto
Temperatura, dia	**22-26 °C**
Ponto quente	**28 °C**
Temperatura, noite	**15-20 °C**
Umidade	**60-80%**
Água	**Sem recipiente de água**
UVB	**Recomendado** Zona de Ferguson 2
Alimentação	**Insetos**
Mantido	**Sozinho**

REPRODUÇÃO

Reprodução	**Ovíparo**
Maturidade sexual	**12 meses**
Tamanho da postura	**2-11 ovos**
Incubação	**180-360 dias** a 18-22 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Quase ameaçado (NT)**

Resumo dos cuidados

O TERRÁRIO	The care of *Kinyongia boehmei* is poorly documented, and several of the figures here are estimated from the climate of the Taita Hills.
TEMPERATURA	By day the temperature should be 22-26 °C with a basking spot of about 28 °C, and it must not rise above 28 °C. At night it should fall to 15-20 °C.
UMIDADE E ÁGUA	Humidity should be 60-80 %. The animal drinks droplets and cannot manage with a water bowl: use a drip or rain system, or mist the leaves several times a day.
ILUMINAÇÃO E UVB	The species is diurnal and needs 12-14 hours of light a day with full-spectrum light and UVB. Ferguson zone 2 is an estimate for a montane chameleon.
ALIMENTAÇÃO	*Kinyongia boehmei* eats only insects. Good feeder insects are small crickets, grasshoppers, houseflies and fruit flies, with mealworms and waxworms now and then.
COMPORTAMENTO E MANUSEIO	*Kinyongia boehmei* is territorial and is stressed by handling and by the sight of conspecifics. Males display with their horns and with colour changes. Do not pick the animal up without reason.
ÁREA DE OCORRÊNCIA NATURAL	Quênia.

O guia completo com mapa e atualizações: herpy.org/en/chameleons/kinyongia-boehmei