herpy

ESCINCÍDEOS

Gold Skink

Eumeces schneiderii

A sturdy, smooth and fast skink from the arid regions between North Africa and Central Asia. *Eumeces schneiderii* is sand-coloured to olive-brown with orange spots, readily burrows into loose sand and is a robust, long-lived terrarium species when given sun, UVB and a deep substrate.

Família: Scincidae Nome anterior: *Eumeces schneideri; Novoeumeces schneideri*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**30-45 cm** comprimento total com a cauda
Longevidade	**15-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre e escavador**
Habitat	**Deserto e semideserto**
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**120 × 60 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**25-30 °C**
Ponto quente	**40-45 °C**
Temperatura, noite	**20-25 °C**
Umidade	**20-40%**
Água	**Recipiente de água**
UVB	**Essencial** Zona de Ferguson 3
Alimentação	**Insetos, frutas e vertebrados**
Mantido	**Sozinho ou um macho com várias fêmeas**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**5-15 ovos**
Incubação	**40-60 dias** a 28-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO

Eumeces schneiderii lives on the ground and burrows, so floor area and substrate are the most important things. A pair, or a male with two females, needs at least 120 × 60 × 45 cm (length × width × height).

TEMPERATURA

The enclosure needs a clear gradient. By day the enclosure should be 25-30 °C, warmest near the basking spot and coolest at the opposite end.

UMIDADE E ÁGUA

The species comes from arid regions and should be kept dry with a humidity of 20-40 %, and in any case below 50 %. Light misting once a week is enough.

ILUMINAÇÃO E UVB

The species is diurnal and needs UVB for as long as the light is on. It belongs to Ferguson zone 3 and needs a UV index of about 3 at the basking spot.

ALIMENTAÇÃO

Eumeces schneiderii is omnivorous, but eats mainly live insects and other small animals: crickets, roaches, mealworms, superworms, silkworms, snails and earthworms.

COMPORTAMENTO E MANUSEIO

The species is curious and readily explores its surroundings. New animals are often cautious, but with patience and feeding from tongs they become calm. Lift the animal from below, and never grab it from above; a bite can hurt.

ÁREA DE OCORRÊNCIA NATURAL

Argélia, Tunísia, Líbia, Egito, Israel, Territórios palestinos, Chipre, Turquia, Síria, Líbano, Jordânia, Irã, Iraque, Arábia Saudita, Geórgia, Armênia, Azerbaijão, Rússia, Cazaquistão, Turcomenistão, Uzbequistão, Tadjiquistão, Quirguistão, Afeganistão e Paquistão.