herpy

ESCINCÍDEOS

Blanchard's Helmet Skink

Tribolonotus blanchardi

Tribolonotus blanchardi is a very small skink from the forests of the Solomon Islands and Bougainville. It measures only 3-4 cm from snout to vent, lives hidden in the damp leaf litter along small forest streams and lays one egg at a time. Little is known about the species and its care.

Família: Scincidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-4 cm** comprimento focinho-cloaca
Longevidade	**10 anos**
Ativo	**De dia e ao crepúsculo**
Estilo de vida	**Terrestre e escavador**
Habitat	**Floresta tropical e floresta de montanha**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**45 × 30 × 30 cm** C × L × A para 1 adulto
Temperatura, dia	**24-26 °C**
Temperatura, noite	**20-22 °C**
Umidade	**80-90%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**1 ovos**
Incubação	**65-85 dias** a 23-27 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Tribolonotus blanchardi*.
TEMPERATURA	*Tribolonotus blanchardi* does not use basking spots. Keep 24-26 °C by day without a basking spot and 20-22 °C at night.
UMIDADE E ÁGUA	*Tribolonotus blanchardi* tolerates drying out very badly. The humidity must be high, 80-90 %, and the substrate must always be damp. Mist morning and evening, or use an automatic rain system.
ILUMINAÇÃO E UVB	*Tribolonotus blanchardi* lives hidden in the leaf litter in shaded forest.
ALIMENTAÇÃO	*Tribolonotus blanchardi* eats insects. Offer small prey: small crickets, fruit flies, springtails, small cockroaches and small earthworms.
COMPORTAMENTO E MANUSEIO	*Tribolonotus blanchardi* is shy and spends most of its time hidden. It is an animal to watch and not for handling. Give a new animal at least two weeks in peace.
ÁREA DE OCORRÊNCIA NATURAL	Ilhas Salomão e Papua-Nova Guiné.