herpy

ESCINCÍDEOS

Spiny Skink

Tribolonotus novaeguineae

Tribolonotus novaeguineae is a small, shy skink from the lowland rainforest of New Guinea with rows of pointed, backward-curving scales down the back. It lives hidden in damp forest litter, is most active at dusk and lays only one egg at a time. Little is known about the care of this particular species, and it is best suited as a display animal.

Família: Scincidae Nome anterior: *Zonurus novaeguineae*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**16-20 cm** comprimento total com a cauda
Longevidade	**10 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Terrestre e escavador**
Habitat	**Floresta tropical e área úmida**
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**60 × 45 × 45 cm** C × L × A para 1 adulto
Temperatura, dia	**24-27 °C**
Ponto quente	**27-28 °C**
Temperatura, noite	**22-24 °C**
Umidade	**70-90%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 1
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Maturidade sexual	**36-48 meses**
Tamanho da postura	**1 ovos**
Posturas por ano	**1-6**
Incubação	**65-85 dias** a 23-27 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Anexo D**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Tribolonotus novaeguineae*. The species lives in the same kind of rainforest floor as *Tribolonotus gracilis* from northern New Guinea, has the same size and largely the same requirements.
TEMPERATURA	The daytime temperature should be 24-27 °C, and a warm hide at 27-28 °C keeps the animal going, even though it does not bask. At night 22-24 °C is suitable.
UMIDADE E ÁGUA	The humidity should be 70-90 %, highest at night, and the substrate must always be damp.
ILUMINAÇÃO E UVB	The species is active at dusk and needs no strong basking spot, but 12 hours of daylight a day maintains its daily rhythm.
ALIMENTAÇÃO	*Tribolonotus novaeguineae* is insectivorous. The staple diet is crickets, cockroaches, earthworms, mealworms and silkworms, and waxworms and superworms are given in smaller amounts.
COMPORTAMENTO E MANUSEIO	*Tribolonotus novaeguineae* is an animal to watch and not for handling. When seized, it may make sounds, play dead, try to flee or drop its tail, and this stresses it.
ÁREA DE OCORRÊNCIA NATURAL	Indonésia e Papua-Nova Guiné.