herpy

GECKOS

Sarasin's giant gecko

Correlophus sarasinorum

Correlophus sarasinorum is a large, rare and nocturnal gecko from the humid forests of the far south of New Caledonia. It resembles a sturdy version of its relative *Correlophus ciliatus*, but has larger eyes, often a pale V-marking on the nape and a far more restricted distribution. The species is vulnerable in the wild and rare in enclosures.

Família: Diplodactylidae Nome anterior: *Rhacodactylus sarasinorum*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**18-28 cm** comprimento total com a cauda
Longevidade	**15-20 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Arborícola**
Habitat	**Floresta tropical**
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**60 × 45 × 90 cm** C × L × A para 1 adulto
Temperatura, dia	**22-26 °C**
Ponto quente	**26-28 °C**
Temperatura, noite	**18-21 °C**
Umidade	**50-80%**
UVB	**Recomendado** Zona de Ferguson 1
Alimentação	**Insetos, frutas e ração completa**
Mantido	**Sozinho**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**1-2 ovos**
Incubação	**60-90 dias** a 22-26 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	There is little documented practice for *Correlophus sarasinorum*, so the figures are based on experience with the smaller *C. ciliatus*.
TEMPERATURA	*Correlophus sarasinorum* needs 22-26 °C by day, and a small area at the top may reach 26-28 °C. At night the temperature should fall to 18-21 °C. Always keep an eye on the heat.
UMIDADE E ÁGUA	The humidity for *Correlophus sarasinorum* should fluctuate as it does in the forest, where nights are damp and days dry out.
ILUMINAÇÃO E UVB	*Correlophus sarasinorum* is active at dusk and at night and sleeps in tree holes by day. How much UVB it needs is not documented.
ALIMENTAÇÃO	There is no published feeding practice for *Correlophus sarasinorum*. It is omnivorous and eats insects and fruit in the wild.
COMPORTAMENTO E MANUSEIO	*Correlophus sarasinorum* is nocturnal and lives hidden. By day it hides, and it becomes properly active only when the light goes off.
ÁREA DE OCORRÊNCIA NATURAL	Nova Caledônia.