herpy

GECKOS

Wiegmann's striped gecko

Gonatodes vittatus

A small, diurnal gecko from Venezuela, Colombia and the islands off the coast, where the male has a white stripe with a black edge down the back and a yellow head. It climbs on tree trunks and walls but cannot climb glass, and it does well in a small enclosure if kept alone or as a pair.

Família: Sphaerodactylidae Nome anterior: *Gymnodactylus vittatus; Gonatodes gillii*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**7-8 cm** comprimento total com a cauda
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta seca**
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**30 × 30 × 30 cm** C × L × A para 2 adultos
Temperatura, dia	**24-27 °C**
Ponto quente	**29-32 °C**
Temperatura, noite	**20-24 °C**
Umidade	**50-60%**
UVB	**Opcional**
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**1 ovos**
Incubação	**65-110 dias** a 25-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair needs an enclosure of at least 30 × 30 × 30 cm (length × width × height), and larger is better, because more space gives more calm.
TEMPERATURA	By day the enclosure should be 24-27 °C. A warm spot of 29-32 °C can be made with a heat mat or a lamp, but it is not necessary if the temperature is otherwise high and stable.
UMIDADE E ÁGUA	*Gonatodes vittatus* thrives in drier air than most tropical geckos. Keep the humidity at 50-60 %.
ILUMINAÇÃO E UVB	The species is diurnal but does not bask, and UVB is optional. Weak UV light can make it easier to get the animals to breed, but too much UV can cause problems with shedding.
ALIMENTAÇÃO	*Gonatodes vittatus* eats insects and other small arthropods.
COMPORTAMENTO E MANUSEIO	The species is bold and curious and is often seen during the day, but it is not an animal to be handled. The skin can tear if the animal is grabbed, and the animals are easily stressed.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia, Venezuela, Guiana, Trinidad e Tobago, Aruba e Curaçao.