herpy

LACERTÍDEOS

Fringe-fingered Lizard

Acanthodactylus erythrurus

Acanthodactylus erythrurus is a fast, slender lizard from Iberia and north-western Africa with long, fringed toes, brown and grey markings and a reddish tail in juveniles and in females during the breeding season. It reaches 18-23 cm, feeds on insects and needs a deep sand substrate, strong sunlight and high heat.

Família: Lacertidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**18-23 cm** comprimento total com a cauda
Longevidade	**5-8 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre e escavador**
Habitat	**Semideserto e bosque e prado temperados**
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**100 × 50 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**25-35 °C**
Ponto quente	**40-45 °C**
Temperatura, noite	**15-22 °C**
Umidade	**30-50%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 3
Repouso de inverno	**10-15 °C** 8-12 semanas
Alimentação	**Insetos**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Maturidade sexual	**10-18 meses**
Tamanho da postura	**1-8 ovos**
Posturas por ano	**1-2**
Incubação	**35-60 dias** a 25-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Acanthodactylus erythrurus*.
TEMPERATURA	*Acanthodactylus erythrurus* is very fond of heat: active adults in the wild maintain a body temperature of 32-38 °C.
UMIDADE E ÁGUA	Humidity should be 30-50 %, as in the dry, open habitat. The sand should be dry, and only the sand in the laying box is kept damp, while the rest of the enclosure is kept well ventilated.
ILUMINAÇÃO E UVB	*Acanthodactylus erythrurus* is diurnal and basks a great deal, so the enclosure needs strong light and UVB.
ALIMENTAÇÃO	*Acanthodactylus erythrurus* eats insects and other small arthropods, in the wild especially ants.
COMPORTAMENTO E MANUSEIO	*Acanthodactylus erythrurus* is fast, wary and shy. It buries itself in the sand when in danger and is hard to catch. Avoid grabbing the animal; the tail is shed under stress. Males are territorial and fight over space.
ÁREA DE OCORRÊNCIA NATURAL	Espanha, Portugal, Gibraltar, Marrocos e Argélia.