herpy

LACERTÍDEOS

Red Tegu

Salvator rufescens

Salvator rufescens is the most heavily built tegu: a diurnal, burrowing lizard from the dry Chaco and western Argentina, Paraguay and Bolivia, which becomes redder with age. Males grow to 140 cm. The species needs a purpose-built enclosure of at least 300 × 150 × 150 cm, a deep burrowing substrate, strong heat and UVB, and it lives for 15-20 years.

Família: Teiidae Nome anterior: *Tupinambis rufescens*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**80-140 cm** comprimento total com a cauda
Longevidade	**15-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre e escavador**
Habitat	**Semideserto e floresta seca**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**300 × 150 × 150 cm** C × L × A para 1 adulto
Temperatura, dia	**24-35 °C**
Ponto quente	**52-57 °C**
Temperatura, noite	**20-23 °C**
Umidade	**70-80%**
Água	**Recipiente de água**
UVB	**Essencial** Zona de Ferguson 3
Repouso de inverno	**15-20 °C** 17-22 semanas
Alimentação	**Insetos, frutas, plantas, vertebrados e minhocas/caracóis**
Mantido	**Sozinho**

REPRODUÇÃO

Reprodução	**Ovíparo**
Maturidade sexual	**24-36 meses**
Tamanho da postura	**15-35 ovos**
Posturas por ano	**1**
Incubação	**40-75 dias** a 29-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little specific knowledge exists about the care of *Salvator rufescens*.
TEMPERATURA	The basking spot should have a surface temperature of 52-57 °C, measured with an infrared thermometer.
UMIDADE E ÁGUA	Air humidity should be 70-80 %, as for the other Argentine tegus. Keep the substrate damp but not muddy by watering it weekly, and mist daily.
ILUMINAÇÃO E UVB	*Salvator rufescens* is diurnal and needs strong light and UVB for 12-14 hours a day.
ALIMENTAÇÃO	*Salvator rufescens* is an omnivore with an emphasis on animal food. Juveniles and animals raised in captivity can be fussy and often prefer meat.
COMPORTAMENTO E MANUSEIO	*Salvator rufescens* has a reputation for being calm and relatively tame and can be trained. Tameness does not come by itself: the animal must be handled calmly and regularly from a young age, and juveniles are shy.
ÁREA DE OCORRÊNCIA NATURAL	Argentina, Paraguai e Bolívia.