herpy

SALAMANDRAS E TRITÕES

## Iberian Ribbed Newt

# *Pleurodeles waltl*

*Pleurodeles waltl* is the largest newt in Europe: a sturdy, flat-headed and almost entirely aquatic species of 15-20 cm with a row of orange warts along the sides, from which the sharp ribs can protrude as a defence. It is hardy, tolerates cold and breeds readily in captivity, but does not thrive in warm water.

Família: Salamandridae  Nome anterior: *Salamandra fenestrata; Salamandra funebris; Salamandra major*

**Toxinas leves na pele**

DADOS PRINCIPAIS

### O ANIMAL

| | |
|---|---|
| Tamanho adulto | **15-20 cm**<br>comprimento total com a cauda |
| Peso | **10-85 g** |
| Longevidade | **10-20 anos** |
| Ativo | **Ao crepúsculo e à noite** |
| Estilo de vida | **Aquático** |
| Habitat | **Área úmida** |
| Toxinas na pele | **Leves**<br>Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca |
| Dificuldade | **Iniciante** |

### CUIDADOS

| | |
|---|---|
| Terrário mín. | **100 × 50 × 50 cm**<br>C × L × A para 5 adultos |
| Temperatura, dia | **18-21 °C** |
| Temperatura, noite | **18-21 °C** |
| Umidade | **60-80%** |
| Água | **Aquaterrário**<br>profundidade 15-20 cm |
| Temperatura da água | **18-21 °C** |
| UVB | **Opcional** |
| Alimentação | **Insetos, ração completa e minhocas/caracóis** |
| Mantido | **Em casal ou em grupo** |

### REPRODUÇÃO

| | |
|---|---|
| Reprodução | **Ovíparo** |
| Desenvolvimento | **Larvas na água** |
| Local de postura | **Na água** |
| Ovos por temporada | **100-1.000** |
| Posturas por ano | **1-2** |
| Eclosão | **7-21 dias**<br>a 20 °C |
| Fase larval | **90-120 dias**<br>da eclosão até a metamorfose |
| Alimentação das larvas | **Outros larvas e pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)** |
| Cuidado parental | **Nenhum** |

### PROTEÇÃO E DOCUMENTAÇÃO

| | |
|---|---|
| CITES | **Não listado** |
| Anexo da UE | **Não listado** |
| IUCN | **Menos preocupante (LC)** |

## Resumo dos cuidados

| | |
|---|---|
| O TERRÁRIO | *Pleurodeles waltl* is almost entirely aquatic and is kept in an aquarium, not in an ordinary terrarium. For a group of five adults, 100 × 50 × 50 cm is a minimum, and more space is better. |
| TEMPERATURA | The water should be 18-21 °C all year round and preferably not above 21 °C. Some animals become stressed by heat, so the tank should stand cool and out of direct sun. |
| UMIDADE E NEBULIZAÇÃO | *P. waltl* is almost entirely aquatic, so the humidity above the water matters little. 60-80 % is plenty, and a tight lid keeps the air humid. The floating land area must be kept wet. Misting is not necessary. |
| ÁGUA E QUALIDADE DA ÁGUA | Outside the breeding season the water is kept 15-20 cm deep. For the breeding season it can be raised. The water must be still and clean with pH 6.9-7.2, ammonia below 0.01 ppm, nitrite below 0.05 ppm and nitrate below 100 ppm. |
| ILUMINAÇÃO E UVB | UVB is not necessary for *P. waltl*, but can be used. Shade and plant cover are more important than strong light, and part of the aquarium must always be in shade. |
| ALIMENTAÇÃO | *P. waltl* is a greedy omnivore that takes most things it can swallow. |
| COMPORTAMENTO E MANUSEIO | *P. waltl* spends almost its whole life in the water and likes to lie hidden under stones and plants. The species is best kept in a group of at least five animals. Adults eat larvae and young, so the young are moved. |
| ÁREA DE OCORRÊNCIA NATURAL | Espanha, Portugal e Marrocos. |