herpy

SALAMANDRAS E TRITÕES

Rough-skinned Newt

Taricha granulosa

Taricha granulosa is a 13-22 cm long newt from the Pacific coast of North America, with a dark, granular back and a bright orange belly. The skin contains tetrodotoxin, one of the most potent toxins found in amphibians, and the animal is deadly poisonous if swallowed. The species alternates between water and land, can be kept in water all year round and must be kept cool.

Família: Salamandridae Nome anterior: *Salamandra granulosa; Triturus granulosus; Triturus similans*

Toxinas fortes na pele

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-22 cm** comprimento total com a cauda
Longevidade	**12-18 anos**
Ativo	**De dia e à noite**
Estilo de vida	**Semiaquático**
Habitat	**Área úmida e bosque e prado temperados**
Toxinas na pele	**Fortes** Evite o contato com a pele e lave bem as mãos depois de mexer no terrário
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**60 × 30 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**18-21 °C**
Temperatura, noite	**14-18 °C**
Umidade	**80-95%**
Água	**Aquaterrário** profundidade 20-30 cm
UVB	**Não necessário**
Alimentação	**Insetos, ração completa e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**Na água**
Maturidade sexual	**48-60 meses**
Ovos por temporada	**7-30**
Eclosão	**21-42 dias**
Fase larval	**60-150 dias** da eclosão até a metamorfose
Alimentação das larvas	**Pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	*Taricha granulosa* is the most aquatic species in the genus. The animals alternate between land and water, some for months or years at a time, but from about one year of age they can be kept in water all year round.
TEMPERATURA	*Taricha granulosa* loves cool conditions. In summer, water and air are kept at 18-21 °C. If it gets warmer, towards 24 °C, the animals move up onto land; prolonged periods above 21 °C should be avoided.
UMIDADE E NEBULIZAÇÃO	The land section and the hides above the water must be damp, but not soaking. Humidity has not been measured for the species; an aquaterrarium with a lid and damp hides gives around 80-95 %.
ÁGUA E QUALIDADE DA ÁGUA	The water must be 20-30 cm deep, and deeper water does no harm: it is almost impossible to give the species water that is too deep in an ordinary tank.
ILUMINAÇÃO E UVB	The species does not need UVB. Ordinary daylight in the room, or a weak fluorescent tube by day and darkness at night, is enough. The tank must not stand in direct sun, and heat lamps must be avoided.
ALIMENTAÇÃO	*Taricha granulosa* is an opportunistic predator. In the wild it eats amphipods, insects, snails, leeches, tadpoles and amphibian and fish eggs.
COMPORTAMENTO E MANUSEIO	Activity varies: some animals are out by day, while others hide under roots and stones and hunt at night. In the breeding season the male holds on to the female in amplexus, which can last from hours to several days.
ÁREA DE OCORRÊNCIA NATURAL	Estados Unidos e Canadá.