herpy

SALAMANDRAS E TRITÕES

California Newt

Taricha torosa

Taricha torosa is a sturdy newt, 13-20 cm long, from the California coast, with a granular brown back and a yellow to orange belly. The skin contains tetrodotoxin, a nerve toxin with no antidote, and the animal can be deadly if swallowed. The species lives mostly on land and enters the water only to breed, and it must be kept cool.

Família: Salamandridae Nome anterior: *Cynops torosus; Notophthalmus torosus* **Toxinas fortes na pele**

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-20 cm** comprimento total com a cauda
Longevidade	**12-22 anos**
Ativo	**De dia e à noite**
Estilo de vida	**Semiaquático**
Habitat	**Área úmida e bosque e prado temperados**
Toxinas na pele	**Fortes** Evite o contato com a pele e lave bem as mãos depois de mexer no terrário
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**Na água**
Maturidade sexual	**36 meses**
Ovos por temporada	**7-30**
Eclosão	**14-21 dias**
Fase larval	**60-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

CUIDADOS

Terrário mín.	**60 × 30 × 30 cm** C × L × A para 2 adultos
Temperatura, dia	**16-21 °C**
Temperatura, noite	**14-18 °C**
Umidade	**70-90%**
Água	**Recipiente de água**
UVB	**Não necessário**
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Quase ameaçado (NT)**

Resumo dos cuidados

O TERRÁRIO	There is relatively little knowledge about the care of *Taricha torosa*, so the winter temperatures are borrowed from the closely related *Taricha granulosa*, which is kept more often.
TEMPERATURA	*Taricha torosa* likes the cool. By day the temperature should be 16-21 °C and at night 14-18 °C. The temperature must not exceed 24 °C, and longer periods above 21 °C must be avoided.
UMIDADE E NEBULIZAÇÃO	The substrate should be damp, but the enclosure should not drip. Humidity is about 70-90%, and daily misting with dechlorinated water keeps the substrate moist.
ÁGUA E QUALIDADE DA ÁGUA	A shallow dish of clean water must always be available, so that the animal can cover itself completely with water. Use tap water treated for chlorine and chloramine; distilled and demineralised water are not used.
ILUMINAÇÃO E UVB	The species does not need UVB. Daylight in the room or a weak fluorescent tube during the day and darkness at night is enough.
ALIMENTAÇÃO	*Taricha torosa* is an opportunistic predator that takes almost anything it can overpower. Earthworms are a complete diet in themselves, and compost worms (*Eisenia fetida*) are also taken readily.
COMPORTAMENTO E MANUSEIO	*Taricha torosa* is calm, sociable and not aggressive towards others of its kind of the same size.
ÁREA DE OCORRÊNCIA NATURAL	Estados Unidos.