herpy

SALAMANDRAS E TRITÕES

Southern Crested Newt

Triturus karelinii

Triturus karelinii is a large, robust crested newt of 12-18 cm from the Crimea, the Caucasus and the southern shore of the Caspian Sea. It has brown to grey, warty skin, and the throat and belly are orange or yellow with small black spots. In spring the male develops a tall, serrated crest. The species is hardy and easy to keep, and it can be kept both on land and in water.

Família: Salamandridae Nome anterior: *Triturus cristatus karelinii* **Toxinas leves na pele**

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**12-18 cm** comprimento total com a cauda
Longevidade	**8-11 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Semiaquático**
Habitat	**Área úmida e bosque e prado temperados**
Toxinas na pele	**Leves** Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca
Dificuldade	**Iniciante**

CUIDADOS

Terrário mín.	**45 × 30 × 30 cm** C × L × A para 4 adultos
Temperatura, dia	**12-20 °C**
Temperatura, noite	**8-15 °C**
Umidade	**70-95%**
Água	**Aquaterrário** profundidade 15-30 cm
Temperatura da água	**10-20 °C**
UVB	**Não necessário**
Repouso de inverno	**4-10 °C** 6-12 semanas
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**Na água**
Maturidade sexual	**12-36 meses**
Ovos por temporada	**150-400**
Eclosão	**14-35 dias**
Fase larval	**56-112 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa, outros larvas e pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	*Triturus karelinii* can be kept following its natural cycle, with an aquatic phase and a terrestrial phase, or in water all year.
TEMPERATURA	The water should be kept below 25 °C, because in warmer water the animals try to leave it. The animals are fully active down to about 10 °C.
UMIDADE E NEBULIZAÇÃO	In the aquatic phase humidity is not an issue. In the terrestrial phase the soft skin dries out quickly, so the substrate must be kept damp, with a humidity of 70-95%.
ÁGUA E QUALIDADE DA ÁGUA	A water depth of 15 cm is enough, and deeper is fine: 15-30 cm gives the animals room to swim. The water is kept at 10-20 °C.
ILUMINAÇÃO E UVB	The animals are nocturnal and do not need UVB or artificial light for their own sake. The plants need light to grow, and an ordinary aquarium lamp is enough.
ALIMENTAÇÃO	In the aquatic phase *T. karelinii* takes a wide range of food: earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms.
COMPORTAMENTO E MANUSEIO	*Triturus karelinii* is robust and easy to keep. In the water the animals snap hard at anything that moves, including each other during feeding, and in the wild crested newts also eat smaller amphibians.
ÁREA DE OCORRÊNCIA NATURAL	Ucrânia, Rússia, Geórgia, Azerbaijão e Irã.