herpy

SALAMANDRAS E TRITÕES

Black Knobby Newt

Tylototriton asperrimus

Tylototriton asperrimus is a dark, warty crocodile newt of 12-15 cm with knobbly rib protuberances and heavy dorsal ridges. The species lives in humid mountain regions in the Chinese province of Guangxi and has given its name to a whole group of closely related crocodile newts, which have only recently been split off as separate species.

Família: Salamandridae Nome anterior: *Echinotriton asperrimus; Pleurodeles asperrimus* **Toxinas leves na pele**

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**12-15 cm** comprimento total com a cauda
Longevidade	**15-25 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Terrestre**
Habitat	**Floresta de montanha e área úmida**
Toxinas na pele	**Leves** Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**60 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**20-24 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Área aquática** profundidade 1-5 cm
UVB	**Opcional**
Repouso de inverno	**8-12 °C** 8-12 semanas
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**No chão**
Ovos por temporada	**30-80**
Eclosão	**20-36 dias** a 22-24 °C
Fase larval	**50-110 dias** da eclosão até a metamorfose
Alimentação das larvas	**Outros larvas e pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Quase ameaçado (NT)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Tylototriton asperrimus*.
TEMPERATURA	In summer the enclosure is kept at 20-24 °C by day and 18-21 °C at night, which corresponds to summer in the range.
UMIDADE E NEBULIZAÇÃO	Humidity is 80-100 %. The enclosure is misted daily with a plant mister, and there must be both damper and drier zones so that the air does not stand still.
ÁGUA E QUALIDADE DA ÁGUA	The water section must have soft, slightly acidic water; rainwater or reverse-osmosis water suits. The pH of the water is not documented for *Tylototriton asperrimus*; *Tylototriton ziegleri* is kept at pH 6.4-6.8.
ILUMINAÇÃO E UVB	Light is of no particular importance to the animals and is arranged for the plants: LED strips or fluorescent tubes for 12 hours in summer and 10-11 hours in winter. UVB is not necessary for this secretive species.
ALIMENTAÇÃO	Adults are given compost worms, crickets, house crickets, waxworms, earthworms and isopods of suitable size about twice a week. In winter they eat irregularly and much less.
COMPORTAMENTO E MANUSEIO	In the closely related species the animals live very secretively and are seen mostly when fed or when the males are in the water. Pairs or small groups can be kept together in a spacious enclosure with many hiding places.
ÁREA DE OCORRÊNCIA NATURAL	China.