herpy

SALAMANDRAS E TRITÕES

Guizhou Crocodile Newt

Tylototriton kweichowensis

Tylototriton kweichowensis is a robust, blackish-brown crocodile newt of 15-21 cm with three broad orange to reddish stripes along the body. It lives on land in the mountains of Guizhou and Yunnan and enters the water mainly to breed. It is best kept in a large enclosure with both land and water.

Família: Salamandridae Nome anterior: *Pleurodeles kweichowensis* **Toxinas leves na pele**

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**15-21 cm** comprimento total com a cauda
Longevidade	**10-20 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Terrestre e semiaquático**
Habitat	**Floresta de montanha e área úmida**
Toxinas na pele	**Leves** Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**90 × 45 × 45 cm** C × L × A para 3 adultos
Temperatura, dia	**18-25 °C**
Temperatura, noite	**15-20 °C**
Umidade	**70-90%**
Água	**Aquaterrário** profundidade 10-15 cm
UVB	**Não necessário**
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**Em folhas sobre a água**
Ovos por temporada	**50-150**
Posturas por ano	**1**
Eclosão	**10-12 dias** a 25 °C
Fase larval	**80-100 dias** da eclosão até a metamorfose
Alimentação das larvas	**Pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	*Tylototriton kweichowensis* needs a large enclosure, because the animals are big and heavy, need a solid land area with a hiding place for each animal, and need enough water for both larvae and a filter.
TEMPERATURA	The species thrives at ordinary room temperature: 18-25 °C by day and 15-20 °C at night. In summer the day may well be at the high end, and in winter the animals are kept at 15-20 °C around the clock.
UMIDADE E NEBULIZAÇÃO	Humidity must be high, but the substrate not wet; 70-90 % is an estimate. Keep the land section damp without letting it turn to mud, and part of the substrate may well be drier.
ÁGUA E QUALIDADE DA ÁGUA	The water section takes up roughly half of the enclosure. The depth is not documented for the species; 10-15 cm is borrowed from *Tylototriton verrucosus*.
ILUMINAÇÃO E UVB	UVB is not necessary, and the species is most active at dusk and at night.
ALIMENTAÇÃO	Adults are fed chopped earthworms, crickets and other prey of suitable size. Waxworms and other fatty larvae are popular, but should not be a staple. With a varied diet, vitamin and mineral supplements are normally not necessary.
COMPORTAMENTO E MANUSEIO	The species is slow and clumsy. 2-3 animals can be kept together when there are enough hiding places for each animal.
ÁREA DE OCORRÊNCIA NATURAL	China.