herpy

SALAMANDRAS E TRITÕES

Wenxian Knobby Newt

Tylototriton wenxianensis

Tylototriton wenxianensis is a medium-sized, almost entirely black crocodile newt with orange toe tips and tail edge. It lives hidden in forest and scrub in the mountains of central China and enters the water only to breed; the eggs are laid on land. Little is known about its care, so this guide is based on close relatives.

Família: Salamandridae **Toxinas leves na pele**

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-16 cm** comprimento total com a cauda
Longevidade	**10-20 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Terrestre e semiaquático**
Habitat	**Floresta de montanha e área úmida**
Toxinas na pele	**Leves** Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**60 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**18-24 °C**
Temperatura, noite	**15-20 °C**
Umidade	**70-90%**
Água	**Área aquática** profundidade 3-5 cm
UVB	**Não necessário**
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Larvas na água**
Local de postura	**No chão**
Ovos por temporada	**30-80**
Eclosão	**20-36 dias** a 22-24 °C
Fase larval	**80-130 dias** da eclosão até a metamorfose
Alimentação das larvas	**Outros larvas e pequenos invertebrados aquáticos (dáfnias, larvas de mosquito)**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Tylototriton wenxianensis*.
TEMPERATURA	In summer the enclosure is kept at 22-24 °C by day, in winter at around 18 °C, and at night it may fall to 15-20 °C. Brief spells of lower temperatures do no harm.
UMIDADE E NEBULIZAÇÃO	In summer the enclosure is misted daily so that the moss and substrate stay moist and the humidity is 70-90 %.
ÁGUA E QUALIDADE DA ÁGUA	The water is soft and slightly acidic, pH 6.4-6.8, and at most about 5 cm deep. Outside the breeding season the animals hardly use the water section; in winter it can be almost dried out, leaving only a little water for humidity.
ILUMINAÇÃO E UVB	UVB is not necessary. The species lives hidden and does not like strong light; the lighting is first and foremost for the plants. Provide about 12 hours of light in summer and 10-11 hours in winter.
ALIMENTAÇÃO	The diet is not documented for the species. The genus takes ordinary newt food: earthworms, crickets, woodlice and waxworms. Adults are fed about twice a week, and in winter they eat irregularly and much less.
COMPORTAMENTO E MANUSEIO	The animals live very secretively and can be kept alone, in pairs or in small groups. Keep the species on its own, without other amphibians. Handle the animals only when necessary, with clean, wet hands or powder-free gloves.
ÁREA DE OCORRÊNCIA NATURAL	China.