herpy

SAPOS E RÃS

Yellow-bellied Toad

Bombina variegata

Bombina variegata is a small, compact frog of 3-6 cm with a grey-brown, sharp-warted back and a bright yellow and blue-grey belly, which it displays when it feels threatened. The species lives in the mountains and hill country of central and southern Europe, favours small, sunny waterholes and is hardy, inquisitive and long-lived in the terrarium. It needs cool temperatures and a cool hibernation, but otherwise little space.

Família: Bombinatoridae

Toxinas leves na pele

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-6 cm** comprimento focinho-cloaca
Peso	**2,3-12 g**
Longevidade	**10-20 anos**
Ativo	**Ao crepúsculo e à noite**
Estilo de vida	**Semiaquático**
Habitat	**Área úmida e bosque e prado temperados**
Toxinas na pele	**Leves** Lave as mãos após o contato e mantenha os dedos longe dos olhos e da boca
Dificuldade	**Iniciante**

CUIDADOS

Terrário mín.	**60 × 30 × 30 cm** C × L × A para 4 adultos
Temperatura, dia	**20-24 °C**
Temperatura, noite	**16-20 °C**
Umidade	**60-80%**
Água	**Área aquática** profundidade 5-10 cm
UVB	**Opcional**
Repouso de inverno	**5-10 °C** 8-12 semanas
Alimentação	**Insetos e minhocas/caracóis**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos na água**
Local de postura	**Na água**
Maturidade sexual	**12-24 meses**
Ovos por postura	**40-170**
Posturas por ano	**1-3**
Eclosão	**2-3 dias**
Fase larval	**25-70 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO
There are not many detailed husbandry accounts of *Bombina variegata*. The recommendations here are based on the species' way of life and on the close relative *Bombina orientalis*, which is far better known in captivity.

TEMPERATURA
The species is adapted to a temperate climate and should not be given heat like most terrarium animals. By day 20-24 °C is suitable, and at night it should be slightly cooler, 16-20 °C.

UMIDADE E NEBULIZAÇÃO
Humidity should be 60-80 %. The large water section holds most of the humidity, so it is normally enough to mist the land section lightly once a day, especially in a dry room in winter.

ÁGUA E QUALIDADE DA ÁGUA
The water section should fill about half the floor area and be 5-10 cm deep.

ILUMINAÇÃO E UVB
Give 12-14 hours of light in summer and down to about 10 hours in winter, controlled by a timer. An LED fixture provides the basic light and makes the plants thrive. No heat lamp is needed. UVB is optional.

ALIMENTAÇÃO
B. variegata has a large appetite and takes anything live that is small enough to fit in the mouth: crickets, small crickets, fruit flies, flies, woodlice, silverfish, small earthworms and small snails.

COMPORTAMENTO E MANUSEIO
B. variegata is inquisitive and sociable and can be kept in pairs or in groups. The animals should be roughly the same size, because large animals can eat small ones, and at feeding they can become a little pushy.

ÁREA DE OCORRÊNCIA NATURAL
França, Bélgica, Países Baixos, Luxemburgo, Alemanha, Suíça, Liechtenstein, Áustria, Itália, Tchéquia, Eslováquia, Polônia, Hungria, Romênia, Moldávia, Ucrânia, Eslovênia, Croácia, Bósnia e Herzegovina, Sérvia, Montenegro, Macedônia do Norte, Albânia, Bulgária e Grécia.