herpy

SAPOS E RÃS

Haraldmeier's Mantella

Mantella haraldmeieri

Mantella haraldmeieri is a small, diurnal frog of 21-27 mm from the south-eastern forests of Madagascar, light brown on the back with black flanks, copper-coloured hind legs and red markings on the underside of the legs. The species is threatened in the wild and only rarely kept, so very little is known about its care. It is kept like the other species in the genus: cool, humid and never warm.

Família: Mantellidae Nome anterior: *Mantella madagascariensis haraldmeieri*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**21-27 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**45 × 45 × 30 cm** C × L × A para 4 adultos
Temperatura, dia	**21-24 °C**
Temperatura, noite	**16-18 °C**
Umidade	**60-80%**
Água	**Recipiente de água**
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos na água**
Local de postura	**No chão**
Maturidade sexual	**12-18 meses**
Ovos por postura	**50-130**
Eclosão	**3-12 dias**
Fase larval	**42-56 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Nenhum**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	Very little is known about keeping *Mantella haraldmeieri*, because the species is only rarely kept.
TEMPERATURA	The temperature should be 21-24 °C during the day and 16-18 °C at night, corresponding to a cool room. Heat should be avoided: over 27 °C can quickly become fatal for the genus. The figures are borrowed from *M. baroni*.
UMIDADE E NEBULIZAÇÃO	Humidity should be 60-70 % day to day and up to 80 % in the breeding season (borrowed from *M. baroni*). Mist the enclosure daily with decalcified water, heavily in warm weather and a little less when it is cool.
ÁGUA E QUALIDADE DA ÁGUA	Place a shallow water dish in the enclosure, and keep it full at all times. The frogs are poor swimmers, so the dish must be so shallow that they can easily get out, and it must not take up much of the floor.
ILUMINAÇÃO E UVB	No UVB requirement is documented for the species. The genus is active in partial sunlight in the wild, so a weak UVB light (Ferguson zone 1) is probably an advantage, provided there is shade and plants so the animals can choose.
ALIMENTAÇÃO	*Mantella haraldmeieri* is fed small arthropods: springtails, wingless fruit flies, newly hatched crickets, small woodlice, aphids and lesser mealworm larvae. The genus takes prey of up to about 6 mm. Variety prevents deficiencies.
COMPORTAMENTO E MANUSEIO	The frog is diurnal and patrols its territory when hunting. The males call, defend territories and may wrestle with one another.
ÁREA DE OCORRÊNCIA NATURAL	Madagascar.