herpy

SAPOS-FLECHA

Brazil-nut Poison Frog

Adelphobates castaneoticus

Adelphobates castaneoticus is a small, glossy black poison frog with white or yellowish spots and bright orange patches at the forelimbs and knees. It lives on the forest floor in the Brazilian state of Pará in the Amazon and is named after the Brazil nut, whose empty shells the male uses as water pools for the tadpoles. Little is known about its care.

Família: Dendrobatidae Nome anterior: *Dendrobates castaneoticus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**18-23 mm** comprimento focinho-cloaca
Longevidade	**8-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**24-27 °C**
Temperatura, noite	**20-24 °C**
Umidade	**70-100%**
Água	**Sem recipiente de água**
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**5-7 meses**
Ovos por postura	**4-8**
Eclosão	**10-14 dias**
Fase larval	**60-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e outros girinos**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Adelphobates castaneoticus*. The species lives in the leaf litter of the forest floor in a warm, humid lowland climate.
TEMPERATURA	The daytime temperature should be 24-27 °C, and at night it falls 3-4 °C, to 20-24 °C. The variation over the year should be small, 1-2 °C. This fits the lowland climate of Pará. There should be no basking spot and no heat lamp.
UMIDADE E NEBULIZAÇÃO	Humidity should be about 70-80 % in the middle of the day and close to 100 % in the morning and evening, and the leaf litter and substrate should be moist at all times.
ÁGUA E QUALIDADE DA ÁGUA	A water dish is not necessary unless you want to raise young. The frogs get their moisture from misting and the moist leaf litter.
ILUMINAÇÃO E UVB	Provide 10-12 hours of light a day with fluorescent tubes or LEDs for the plants. In the wild the species lives in the shade under the forest canopy, so the light should give the plants enough to grow without heating the enclosure.
ALIMENTAÇÃO	In the wild *A. castaneoticus* feeds on ants, termites and other small arthropods. In captivity feed springtails, small fruit flies and small isopods.
COMPORTAMENTO E MANUSEIO	*A. castaneoticus* is diurnal and lives mostly on the forest floor. Temperament and territorial behaviour are not documented, but *A. galactonotus* can become aggressive towards conspecifics when it becomes sexually mature.
ÁREA DE OCORRÊNCIA NATURAL	Brasil.