herpy

SAPOS-FLECHA

Rio Madeira Poison Frog

Adelphobates quinquevittatus

Adelphobates quinquevittatus is a very small, glossy black poison frog of 14-16 mm with white stripes and grey, greenish or orange legs. It lives in the leaf litter of the rainforest in the Madeira River drainage in the southern Amazon and is only rarely kept. Very little is known about its care.

Família: Dendrobatidae Nome anterior: *Dendrobates quinquevittatus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**14-16 mm** comprimento focinho-cloaca
Longevidade	**8-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**24-27 °C**
Temperatura, noite	**20-24 °C**
Umidade	**70-100%**
Água	**Sem recipiente de água**
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-8**
Eclosão	**10-14 dias**
Fase larval	**60-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e outros girinos**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Very little is known about the care of *Adelphobates quinquevittatus*. The species lives in the leaf litter in a warm and humid lowland climate.
TEMPERATURA	The daytime temperature should be 24-27 °C, and at night it falls 3-4 °C, to 20-24 °C. The variation over the year should be small, 1-2 °C. This fits the lowland climate of Rondônia.
UMIDADE E NEBULIZAÇÃO	Humidity should be about 70-80 % in the middle of the day and close to 100 % in the morning and evening, and the leaf litter and substrate should be moist at all times.
ÁGUA E QUALIDADE DA ÁGUA	A water dish is not necessary unless you want to raise young. The frogs get their moisture from misting and the moist leaf litter.
ILUMINAÇÃO E UVB	Provide 10-12 hours of light a day with fluorescent tubes or LEDs for the plants. In the wild the species lives in the shade under the forest canopy, so the light should give the plants enough to grow without heating the enclosure.
ALIMENTAÇÃO	In the wild *A. quinquevittatus* feeds mostly on ants and termites. In captivity feed small fruit flies (*Drosophila melanogaster*), springtails, micro crickets, small waxworm larvae, sieved meadow plankton and small isopods.
COMPORTAMENTO E MANUSEIO	*A. quinquevittatus* is diurnal and lives mostly in the leaf litter. Temperament and territorial behaviour are not documented.
ÁREA DE OCORRÊNCIA NATURAL	Brasil.

O guia completo com mapa e atualizações: herpy.org/en/poison-dart-frogs/adelphobates-quinquevittatus