herpy

SAPOS-FLECHA

Sanguine Poison Frog

Allobates zaparo

Allobates zaparo is a small, diurnal frog of about 3 cm from the rainforest at the foot of the Andes in Ecuador and northern Peru. Adults have a red back, black flanks, a blue belly and yellow spots at the limbs and mimic the poisonous *Ameerega bilinguis* and *Ameerega parvula*, but are not poisonous themselves. The species is only rarely kept, so the care is based on the climate and on its relative *Allobates femoralis*.

Família: Aromobatidae Nome anterior: *Epipedobates zaparo; Phyllobates zaparo; Dendrobates zaparo*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3 cm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele	**Nenhuma conhecida**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**60 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**24-27 °C**
Temperatura, noite	**21-23 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**8-20**
Eclosão	**15-20 dias**
Fase larval	**40-50 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Allobates zaparo*. The species is only rarely kept, and captive breeding is not described.
TEMPERATURA	Keep the terrarium at 24-27 °C by day and 21-23 °C at night. The figures follow the climate of Ecuador's lowlands at the foot of the Andes. The variation over the year should be small, 1-2 °C. There is no winter rest.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-80 % in the middle of the day and close to 100 % in the morning and evening.
ÁGUA E QUALIDADE DA ÁGUA	Provide a shallow glass dish with 1-2 cm of water that the frogs can easily climb out of. The dish is used by the adults when the tadpoles are to be deposited.
ILUMINAÇÃO E UVB	The light is first and foremost for the plants: 10-12 hours a day with LED light with a plant-suitable spectrum. It is not documented that *Allobates zaparo* needs UVB.
ALIMENTAÇÃO	In the wild ants are an important part of the diet.
COMPORTAMENTO E MANUSEIO	*Allobates zaparo* is diurnal and lives in the leaf litter. Males call from the top of the leaves to attract females.
ÁREA DE OCORRÊNCIA NATURAL	Equador e Peru.