herpy

SAPOS-FLECHA

Pleasing Poison Frog

Ameerega bassleri

Ameerega bassleri is a diurnal poison frog of just under 4 cm from the humid forests at the eastern foot of the Andes in northern Peru. The male guards the eggs and carries the tadpoles to the water, and the species must be kept cooler than most poison frogs. Little is known about its care, and the species is vulnerable in the wild.

Família: Dendrobatidae Nome anterior: *Epipedobates bassleri; Dendrobates bassleri; Phobobates bassleri*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**4 cm** comprimento focinho-cloaca
Longevidade	**8-12 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**50 × 50 × 55 cm** C × L × A para 2 adultos
Temperatura, dia	**18-21 °C**
Temperatura, noite	**13-17 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Maturidade sexual	**12-13 meses**
Ovos por postura	**40**
Eclosão	**10 dias**
Fase larval	**50-60 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Ameerega bassleri*. The species lives in humid forest by streams at 300-1,200 m, where the air is cooler than for most poison frogs.
TEMPERATURA	The species must not be kept too warm. By day 18-21 °C suits, and at night the temperature should fall 4-6 °C, to about 13-17 °C. The difference between the seasons is only 1-2 °C.
UMIDADE E NEBULIZAÇÃO	Humidity must be 70-100 %. Morning and evening it is close to 100 % with mist, while around midday it may fall to about 70 %.
ÁGUA E QUALIDADE DA ÁGUA	A water bowl with 1-2 cm of water must be in the vivarium, and a permanent water area or small containers are used for the tadpoles when breeding. Use rainwater or reverse osmosis water and change the water several times a week.
ILUMINAÇÃO E UVB	The species lives on the forest floor, where little light reaches. LED light with a plant-suitable spectrum and a 10-12 hour day length is enough.
ALIMENTAÇÃO	The species takes the usual small feeder insects: fruit flies, springtails, newly hatched crickets, the smallest waxworm larvae and sweep-netted small insects from the meadow.
COMPORTAMENTO E MANUSEIO	The species is diurnal and lives on the ground. The male is territorial and calls to defend his area, so the animals need many hiding places and broken lines of sight.
ÁREA DE OCORRÊNCIA NATURAL	Peru.