herpy

SAPOS-FLECHA

Yurimaguas Poison Frog

Ameerega hahneli

Ameerega hahneli is a small, diurnal poison frog of 17-22 mm with pale side stripes and a blue belly, which lives in the rainforest leaf litter across large parts of the Amazon lowlands. The male carries the tadpoles to running water, and the tadpoles need current in the water. The species is common in the wild but rare in captivity.

Família: Dendrobatidae Nome anterior: *Epipedobates hahneli; Dendrobates hahneli*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**17-22 mm** comprimento focinho-cloaca
Longevidade	**6-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**7-12 meses**
Ovos por postura	**6-33**
Eclosão	**4-16 dias**
Fase larval	**55-65 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**50 × 50 × 50 cm** C × L × A para 4 adultos
Temperatura, dia	**23-25 °C**
Temperatura, noite	**19-21 °C**
Umidade	**65-90%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Only limited knowledge exists about the care of *Ameerega hahneli*, and the requirements may vary between the different forms and possible hidden species.
TEMPERATURA	*Ameerega hahneli* needs 23-25 °C by day and 19-21 °C at night. Fluorescent tubes outside the terrarium can supply both light and the necessary warmth. Above 28 °C there is a risk of stress; cool with a fan and extra spraying.
UMIDADE E NEBULIZAÇÃO	Humidity must be 65-90 %.
ÁGUA E QUALIDADE DA ÁGUA	There must always be a little water to go to, for example in the drainage channel or a shallow water bowl with 1-2 cm of water. In the dry season the channel is not emptied, so the animals still have access to water.
ILUMINAÇÃO E UVB	The light can be fluorescent tubes or LED with a plant-suitable spectrum for 10-12 hours. UVB is not necessary; a weak UVB bulb (Ferguson zone 1) can be used when there are hides so that the animals can withdraw from the light.
ALIMENTAÇÃO	The small fruit fly (*Drosophila melanogaster*) is the most important food. Adults also take the large fruit fly (*Drosophila hydei*), but often skip it because it is a little too big.
COMPORTAMENTO E MANUSEIO	The species is diurnal and lives on the ground, but is most active early in the morning and late in the afternoon and evening; the rest of the day it is shyer.
ÁREA DE OCORRÊNCIA NATURAL	Brasil, Bolívia, Peru, Equador, Colômbia, Venezuela, Guiana, Guiana Francesa e Suriname.