herpy

SAPOS-FLECHA

Ruby Poison Frog

Ameerega parvula

Ameerega parvula is a small, diurnal poison frog of 18-24 mm from the rainforest leaf litter in the western Amazon lowlands and at the foot of the Andes. The back is black with dark red spots, and the underside is marbled blue and black. The species is rarely kept, and little is known about its care.

Família: Dendrobatidae Nome anterior: *Epipedobates parvulus; Dendrobates parvulus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**18-24 mm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**25-26 °C**
Temperatura, noite	**20-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-16**
Eclosão	**10-22 dias**
Fase larval	**70-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Ameerega parvula*.
TEMPERATURA	The daytime temperature is 25-26 °C, and at night it is lowered by 4-5 degrees to 20-22 °C.
UMIDADE E NEBULIZAÇÃO	Humidity is 70-80 % by day, with a dip to about 70 % around midday, and close to 100 % morning and evening, preferably with mist.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water bowl with 1-2 cm of water lets the frogs drink and is necessary if breeding is to take place, because the male deposits the tadpoles there. The bowl must have a slope or stones so that a frog can climb out again.
ILUMINAÇÃO E UVB	The light should first and foremost give the plants a day-night rhythm of about 12 hours, and an LED plant lamp is enough for that. UVB is not necessary for poison frogs, so it is optional.
ALIMENTAÇÃO	The food is small, live arthropods: springtails and fruit flies (*Drosophila melanogaster*), micro crickets, sieved meadow plankton and small isopods.
COMPORTAMENTO E MANUSEIO	The species is diurnal and lives mostly on the ground. The males call and probably defend small territories, as their relatives do, so a pair is the safe way to keep them in a smaller vivarium.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia, Equador e Peru.