herpy

SAPOS-FLECHA

Peru Poison Frog

Ameerega petersi

Ameerega petersi is a small, diurnal poison frog of 24-30 mm from the rainforest at the foot of the Andes in eastern Peru and western Brazil. The back is brown with a pale, oblique side stripe. The species is rarely kept, and little is known about its care.

Família: Dendrobatidae Nome anterior: *Epipedobates petersi; Dendrobates petersi*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**2-3 cm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**25-28 °C**
Temperatura, noite	**21-25 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-16**
Eclosão	**10-22 dias**
Fase larval	**70-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Ameerega petersi*, but the species is kept in small numbers, and the basic figures for the vivarium are known. It is kept like other small *Ameerega* species from the rainforest leaf litter.
TEMPERATURA	The daytime temperature is 25-28 °C, and at night it is lowered by 3-4 degrees to 21-25 °C. The variation over the year is only 1-2 degrees.
UMIDADE E NEBULIZAÇÃO	Humidity must be 70-100 %: close to 100 % morning and evening, preferably with mist, and about 70-80 % around midday. Mist with reverse osmosis or rainwater morning and evening, or use a fogger or rain system on a timer.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water bowl with 1-2 cm of water is enough. The bowl must be shallow enough for a froglet to climb out of it, and have stones or moss at the edge. Change the water often, and clean the bowl with hot water without soap.
ILUMINAÇÃO E UVB	Provide an LED light with a plant-suitable spectrum for 10-12 hours a day. UVB is not necessary for poison frogs; a weak UVB light (Ferguson zone 1) is optional.
ALIMENTAÇÃO	The diet consists of small live arthropods from the leaf litter.
COMPORTAMENTO E MANUSEIO	The males in the genus call and defend small territories. How *Ameerega petersi* behaves in groups has not been described, so a pair is the safest way to keep them.
ÁREA DE OCORRÊNCIA NATURAL	Peru e Brasil.