herpy

SAPOS-FLECHA

Cordillera Azul Poison Frog

Ameerega rubriventris

Ameerega rubriventris is a small, dark poison dart frog of 21-25 mm with pale side stripes and a red-black or orange-black marbled belly. It lives only in the forests of the Cordillera Azul in central Peru and is threatened. The species is rarely kept, and little is known about its care.

Família: Dendrobatidae Nome anterior: *Epipedobates rubriventris*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**21-25 mm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-16**
Eclosão	**10-22 dias**
Fase larval	**70-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**50 × 50 × 50 cm** C × L × A para 2 adultos
Temperatura, dia	**25-28 °C**
Temperatura, noite	**21-25 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Ameerega rubriventris*.
TEMPERATURA	The daytime temperature is 25-28 °C, and at night it is lowered by 3-4 degrees to 21-25 °C. The temperature varies only 1-2 degrees over the year, and a winter rest period is not necessary.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-80 % by day and close to 100 % morning and evening, ideally with mist. A rain system or a misting machine on a timer makes this easy; otherwise spray thoroughly morning and evening.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water bowl with 1-2 cm of water is enough, and it should be easy to get in and out of.
ILUMINAÇÃO E UVB	The light should primarily serve the plants: an LED plant lamp for 10-12 hours a day and a clear night. UVB is not necessary for poison dart frogs; a weak UVB lamp (Ferguson zone 1) is optional.
ALIMENTAÇÃO	The frogs eat small live arthropods: fruit flies (*Drosophila melanogaster* and, for adults, also *D. hydei*), springtails, micro crickets, small isopods and sieved meadow plankton.
COMPORTAMENTO E MANUSEIO	*Ameerega rubriventris* is diurnal and stays mainly on the forest floor, but readily climbs. The male calls with a bird-like peeping, most often 2-3 short whistles in a row.
ÁREA DE OCORRÊNCIA NATURAL	Peru.