herpy

SAPOS-FLECHA

Three-striped Poison Frog

Ameerega trivittata

Ameerega trivittata is a large, diurnal poison frog of 4-6 cm from the forest floor of the Amazon. It is black with two pale green stripes along the flanks, and in a single trip the male often carries 15-30 tadpoles on his back to water. The frog needs a more spacious enclosure than the small poison frogs.

Família: Dendrobatidae Nome anterior: *Epipedobates trivittatus; Dendrobates trivittatus; Phobobates trivittatus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**4-6 cm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e área úmida**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**12-24 meses**
Ovos por postura	**15-40**
Eclosão	**15-22 dias**
Fase larval	**40-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**70 × 50 × 50 cm** C × L × A para 2 adultos
Temperatura, dia	**23-25 °C**
Temperatura, noite	**19-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	An enclosure of at least 70 × 50 × 50 cm (length, width, height) suits two adult *Ameerega trivittata*.
TEMPERATURA	Keep the daytime temperature at 23-25 °C, and lower it at night by 3-4 °C to about 19-22 °C. The variation over the year should be small, 1-2 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-90 % for most of the day, close to 100 % morning and evening, and it may fall to about 70 % in the middle of the day. A longer time below 70 % is dangerous, especially if the frog has no access to water.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish with 1-2 cm of water is enough. The dish must be shallow enough for a young frog to climb out, with stones or moss at the edge.
ILUMINAÇÃO E UVB	UVB is not documented as necessary for poison frogs in general, and for this particular species there is no information. A weak UVB light (Ferguson zone 1) is optional.
ALIMENTAÇÃO	The diet consists of small live arthropods.
COMPORTAMENTO E MANUSEIO	The males call loudly and defend small territories, while the females are calm. It is not documented how the species behaves in groups in captivity, so pairs are the safest way to keep them.
ÁREA DE OCORRÊNCIA NATURAL	Bolívia, Brasil, Colômbia, Guiana, Peru, Suriname e Venezuela.