herpy

SAPOS-FLECHA

Cauca Poison Frog

Andinobates bombetes

Andinobates bombetes is a small, diurnal poison dart frog of 17-19 mm from the cloud forests of Colombia's Andes. It is black to dark brown with red or orange markings on the front of the body and depends on water-filled bromeliads. The species is Vulnerable in the wild, and little is known about its care.

Família: Dendrobatidae Nome anterior: *Ranitomeya bombetes; Dendrobates bombetes; Minyobates bombetes*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**17-19 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**1-2**
Eclosão	**16-24 dias**
Fase larval	**108-112 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

CUIDADOS

Terrário mín.	**45 × 50 × 60 cm** C × L × A para 5 adultos
Temperatura, dia	**22-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Sem recipiente de água**
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Andinobates bombetes*. What is known is based on the habitat, a cool and very humid mountain forest with thick leaf litter and bromeliads, and on captive breeding in Colombia.
TEMPERATURA	The daytime temperature should be 22-25 °C, and at night it may fall to 18-21 °C as in the mountain forest. Keep the temperature below 27 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 % and never below 70 %. Spray twice a day, morning and afternoon, or use a fogger or rain system with reverse osmosis water or rainwater.
ÁGUA E QUALIDADE DA ÁGUA	A water bowl is not needed; the frogs drink droplets from plants and spray water. For the tadpoles, use bromeliads with water in the leaf axils or small plastic cups with 2-3 cm of water.
ILUMINAÇÃO E UVB	The light is primarily for the plants: plant lighting for 10-12 hours a day provides a daily rhythm and growth.
ALIMENTAÇÃO	The food is small arthropods: springtails, small fruit flies (*Drosophila melanogaster*), small isopods and newly hatched crickets. Newly metamorphosed froglets take only springtails.
COMPORTAMENTO E MANUSEIO	*Andinobates bombetes* is diurnal and stays low in the enclosure, in the leaf litter and by the bromeliads. The males call mainly in the morning.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.